arXiv:1208.5876v1 [math.CA] 29 Aug 2012

A sharp L_p-L_q-Fourier restriction theorem for a conical surface of finite type

Stefan Buschenhenke

Christian-Albrechts-Universität zu Kiel

30th August 2012

Contents

1 Introduction and first considerations 2

1.1 Introduction . . . 2

1.2 The main theorem and necessary conditions . . . 3

1.3 Reduction . . . 5

2 Reduction of the problem 7

2.1 Decomposition . . . 7

2.2 Proceeding to the crux of the problem . . . 9

3 Dense families 12

3.1 Basic properties . . . 12

3.2 An application . . . 13

4 Estimation of the overlap 17

4.1 Straightforward results . . . 17

4.2 Further results . . . 18

4.3 Summary of the results . . . 25

5 L_p-estimates for convolutions 28

5.1 Introduction to the Problem and formulation of the result . . . 28

5.2 Proof . . . 29

6 Proof of theorem 2.2 32

6.1 Application of the results from chapter 5 . . . 32

6.2 Completing the proof . . . 35

7 Appendix: Auxiliary lemmas 40

7.1 Classical lemmas . . . 40

7.2 More special lemmas . . . 41

References 44

1 Introduction and first considerations

1.1 Introduction

Let S be a compact hypersurface in $\mathbb{R}^n$ (or more general a smooth submanifold) with surface measure σ. We say that the $L_p(\mathbb{R}^n)$-$L_q(S)$ *Restriction Theorem* holds if there exists a constant $C > 0$ such that

$$\left(\int\limits_S |\hat{f}(\xi)|^q \, \mathrm{d}\sigma(\xi) \right)^{\frac{1}{q}} \leq C \cdot \|f\|_p \tag{1}$$

holds for every Schwartz function f. A good overview of this topic offers the famous book of E. M. Stein [St]. More detailled information concerning the development in this research field may be found in the overview article of Terence Tao [T].
The object of my investigations is a surface of so-called finite typ, where the tangent plane has finite order of contact. This means - describing the surface locally as a graph of a function - that the second derivative of the function with respect to some direction vanishes at some points (and therefore the curvature as well). One just demands not all derivatives to vanish. A simple example is the curve x^m, $m \geq 3$. However, the restriction estimates for curves are known since 1979 [ACK]. But we can construct a cone-like surface from the curve, like the „classical“ cone is arised from a circle. A sharp restriction theorem for the „classical“ cone was proven by Barcelo [Ba1], remarkably without bilinear estimates. The reason for this is that the cone somehow behaves like a curve, and in the one-dimensional situation, you may use other methods.
Unfortunatly, just when I finished my work, I was informed that Barcelo himself already dealt with the same problem in another paper [Ba2], which was unknown to me. However, although the basic structur might be comparable at the end, the details and techniques are quite different. This can be seen clearest in the different choice of decompositions, which reveal the discrepancy between Barcelos point of view and my own.
Another point is that Barcelo always considered the full cone (equipped with a modified measure to compensate scaling) and not just a compact part. But, at least away from a sharp line, both approaches are equivalent. Furthermore, as the „conical direction“ gives neither curvature nore a finite type condition, one would expect no impact from here, and my notation allows to verify this. Besides obtaining a similar result independently, the most striking point certainly is that unlike Barcelo, I was able to get the sharp range of p and q.
There is a technical obstacle, preventing Barcelo from obtaining the optimal result. When I tried to adopt Barcelos arguments, I got into trouble with the same obstacle quickly. But eventually, I discovered a way to outmaneuver it. The „correct“ formulation of the problem I found, and which I could prove, can be seen in terms of the affine arclength measure, which is constructed in order to compensate the decreasing curvature. Such variations of restriction theorems can be found in Drury and Marshall [DM], who also considered curves of finite typ. This is of course a different problem. But like in [DM], we can derive the original restriction estimate from the estimate with affine arclength measure, with a different range of p and q.
The development of my proof might be of interest for the reader in so far, as I first proved

the (already well-known) result for curves and secondly adopted it for the generalised cone. Hence, the ideas and pictures will often be just in x-$y-$space.

Following the exact formulation of the problem and reduction on the essential core of it, in the second part the mentioned step with the affine arclenght measure will be explained. A further reduction will be achieved by a certain approximation, simultanously both of the function and of the surface measure, like in [Ba1]. The precise arguments are given in the following chapter. For this approximation we decompose the surface into special pieces on each of which the curvature will be almost constant. Furthermore, we need to estimate the overlap of the sets we obtain by taking the sum of two sets of the original decomposition. In Chapter 5, we want to investigate on the convolution of the characteristic functions (or smooth versions) of two sets of our decomposition.
Well prepared by these instruments, we are ready to conclude the proof. The last chapter containes some technical details and auxiliary lemmas.

1.2 The main theorem and necessary conditions

Let γ be a compact curve of finite typ, i.e. for all $p \in \gamma$ exists a local parametrisation Φ, $\Phi(x) = p$, and $m = m_p \in \mathbb{N}_{\geq 2}$ such that Φ is $m-$times differentiable at x and $\Phi^{(m)}(x) \neq 0$. By compactness, we may assume that there exist a universale M such that we allways can find such a m_p bounded by M. Moreover, we choose M minimal with this property.

Theorem 1.1
Let $1 \leq p, q \leq \infty$. *We consider the generalised cone*
$\Gamma = \{(\xi, z) \in \mathbb{R}^2 \times \mathbb{R} |\, 1 \leq z \leq 2,\ \frac{\xi}{z} \in \gamma\}$ *with surface measure* σ. *Then the Fourier restriction theorem*

$$\|\hat{f}|_\Gamma\|_{L_q(\Gamma,\sigma)} \leq C\|f\|_{L_p(\mathbb{R}^3)} \qquad \forall f \in \mathcal{S}(\mathbb{R}^3), \tag{2}$$

holds if $1 \leq p < \frac{M+1}{M}$ *and* $\frac{1}{q} \geq \frac{M+1}{p'}$.

The theorem is sharp in the sense that there exists surfaces where (2) is not valid if the conditions on p' and q are hurt. First we deal with this necessity of the conditions. Instead of doing so in full generality, To convince us of this necessity, we will discuss an example in $\mathbb{R}^2$, and for simplicity omit the cone. This might be more helpfull in order to just understand the idea behind it. We consider the curve $\gamma = \{(x, x^m) | x \in [0, 1]\}$ in $\mathbb{R}^2$: Assume the restriction estimate

$$\|\hat{f}|_\gamma\|_{L_q(\gamma,\sigma)} \leq C\|f\|_{L_p(\mathbb{R}^2)} \qquad \forall f \in \mathcal{S}(\mathbb{R}^2) \tag{3}$$

holds for some $q > \frac{p'}{m+1}$. The following is a variation of the classical Knapp Box example. We define the anisotropic dilatation of a function $f \in \mathcal{S}$ with respect to $r > 0$ by $f_r(x, y) =$

$f(rx, r^m y)$, such that $\hat{f}_r(\xi,\eta) = r^{-(m+1)}\hat{f}\left(\frac{\xi}{r}, \frac{\eta}{r^m}\right)$. We apply (3) for f_r, $r \leq 1$: On the left-hand-side we get

$$\begin{aligned}\|\hat{f}_r\|_{L_q(\gamma,\sigma)} &= \left(\int_0^1 |r^{-(m+1)}\hat{f}\left(\tfrac{\xi}{r}, \tfrac{\xi^m}{r^m}\right)|^q \,\mathrm{d}\xi\right)^{\frac{1}{q}} \\ &= r^{-(m+1)}\left(\int_0^{\frac{1}{r}} |\hat{f}\left(\xi,\xi^m\right)|^q r \,\mathrm{d}\xi\right)^{\frac{1}{q}} \\ &\underset{r\leq 1}{\geq} r^{\frac{1}{q}-(m+1)}\left(\int_0^1 |\hat{f}\left(\xi,\xi^m\right)|^q \,\mathrm{d}\xi\right)^{\frac{1}{q}} \\ &= r^{\frac{1}{q}-(m+1)}\|\hat{f}\|_{L_q(\gamma,\sigma)},\end{aligned}$$

and therefore

$$r^{\frac{1}{q}-(m+1)}\|\hat{f}\|_{L_q(\gamma,\sigma)} \leq \|\hat{f}_r\|_{L_q(\gamma,\sigma)} \leq C\|f_r\|_p = C\,r^{-\frac{1+m}{p}}\|f\|_p.$$

We conclude that

$$r^{\frac{1}{q}-(m+1)+\frac{1+m}{p}}\|\hat{f}\|_{L_q(\gamma,\sigma)} \leq C\|f\|_p,$$

where the exponent of r is negativ: $\frac{1}{q}-(m+1)+\frac{1+m}{p} = \frac{1}{q}-(m+1)\left(1-\frac{1}{p}\right) = \frac{1}{q}-\frac{m+1}{p'} < 0$. We get a contradiction by choosing r small enough, and therefore $\frac{1}{q} \geq \frac{m+1}{p'}$. Since $q \geq 1$, we may already conclude that $p' \geq m+1$. That the case of equality is excluded can be proven by results of the theory of oscillatory integrals:
Again consider $\gamma = \{(x, x^m) | x \in [0,1]\}$ and assume that (3) and respectively

$$\|\widehat{f\,\mathrm{d}\sigma}\|_{L_{p'}(\mathbb{R}^2)} \lesssim \|f\|_{L_{q'}(\gamma)} \tag{4}$$

holds for $p' = m+1$. It is known that [Bu]

$$I(\lambda) = \int e^{-i(x^m+\lambda x)}\,\mathrm{d}x \approx \lambda^{-\frac{m-2}{2m-2}} \quad \text{as } \lambda \to +\infty. \tag{5}$$

We obtain for $f \equiv 1$

$$\begin{aligned}\widehat{f\,\mathrm{d}\sigma}(a,b) &\approx \int_0^1 e^{-i(ax+bx^m)}\,\mathrm{d}x \\ &= b^{-\frac{1}{m}}\int_0^{b^{\frac{1}{m}}} e^{-i(ax/b^{\frac{1}{m}}+x^m)}\,\mathrm{d}x \\ &\overset{(5)}{\approx} b^{-\frac{1}{m}}\left(\frac{a}{b^{\frac{1}{m}}}\right)^{-\frac{m-2}{2m-2}} \qquad \text{if } b \gg 1 \\ &= a^{-\frac{m-2}{2m-2}}b^{-\frac{1}{2m-2}}, \qquad \text{if } \left|\frac{a}{b^{\frac{1}{m}}}\right| \gg 1.\end{aligned}$$

and therefore in the case $m > 3$, i.e. $1 > \frac{m+1}{2m-2} = \frac{p'}{2m-2}$

$$\begin{aligned}
\|\widehat{f\ \mathrm{d}\sigma}\|_{p'}^{p'} &\geq \int_1^\infty \int_C^{Ca^m} |\widehat{f\ \mathrm{d}\sigma}|^{p'}\ \mathrm{d}b\ \mathrm{d}a \\
&\approx \int_1^\infty a^{-p'\frac{m-2}{2m-2}} \int_1^{a^m} b^{-\frac{p'}{2m-2}}\ \mathrm{d}b\ \mathrm{d}a \\
&\approx \int_1^\infty a^{-p'\frac{m-2}{2m-2}} \left[a^{m(1-\frac{p'}{2m-2})} - 1\right]\ \mathrm{d}a \\
&\approx \int_1^\infty a^{-p'\frac{m-2}{2m-2}} a^{m(1-\frac{p'}{2m-2})}\ \mathrm{d}a \\
&= \int_1^\infty a^{m-p'}\ \mathrm{d}a.
\end{aligned}$$

This integral converges if and only if $m-p' < 1$, i.e. $p' > m+1$ or, equivalently, if $p < \frac{m+1}{m}$. The case $m = 3$, which means $1 = \frac{m+1}{2m-2} = \frac{p'}{2m-2}$ and $p'\frac{m-2}{2m-2} = 1$, yields

$$\begin{aligned}
\|\widehat{f\ \mathrm{d}\sigma}\|_{p'}^{p'} &\geq \int_1^\infty \int_C^{Ca^m} |\widehat{f\ \mathrm{d}\sigma}|^{p'}\ \mathrm{d}b\ \mathrm{d}a \\
&\approx \int_1^\infty a^{-p'\frac{m-2}{2m-2}} \int_1^{a^m} b^{-\frac{p'}{2m-2}}\ \mathrm{d}b\ \mathrm{d}a \\
&\approx \int_1^\infty a^{-1} \log a\ \mathrm{d}a \\
&\gtrsim \int_1^\infty a^{-1}\ \mathrm{d}a = \infty.
\end{aligned}$$

However, even in the case $p' = m+1$ a weak-typ estimate might still hold. This is beyond the methods of this paper.

1.3 Reduction

Let γ be a compact curve of finite typ and M the maximal order of contact like above. Obviously, it suffice to decompose γ into finitely many parts and prove the theorem for each part. At first, we cut the curve into finitely many parts, where the curve (after a permutation of coordinates, if necessary) can be described as the graph of some function $\Phi : [a,b] \to \mathbb{R}$. Then we decompose each such part again: For every $x_0 \in [a,b]$ with $\Phi''(x_0) = 0$, we will prove the restriction theorem for $\Phi|_{[x_0-\varepsilon, x_0+\varepsilon]}$ for some $\varepsilon > 0$ (depending on x_0). The compact set $\{\Phi'' = 0\}$ is covered by finitely many of these intervalls. But outside an arbitrary open neighbourhood of the set $\{\Phi'' = 0\}$, the curvature is bounded from above and from below, and in this case results are known [Ba1]. So, how to determine ε?

Given $\Phi''(x_0) = 0$, choose $m \geq 2$ with $\Phi^{(m)}(x_0) \neq 0$ minimal, thus $m \geq 3$ and $\Phi''(x_0) = \ldots = \Phi^{(m-1)}(x_0) = 0$. We claim that we may assume

$$x_0 = 0 \qquad \text{and} \qquad \Phi(0) = \Phi'(0) = 0. \tag{6}$$

For this purpose we introduce the affine transformation $\tilde{\Phi}(x - x_0) = \Phi(x) - \Phi(x_0) - (x - x_0)\Phi'(x_0)$, which provides $\tilde{\Phi}(0) = 0 = \tilde{\Phi}'(0)$, whereas $\tilde{\Phi}^{(k)}(x-x_0) = \Phi^{(k)}(x)$ (and therefore

leaving the finite typ condition untouched).
If we define $\gamma_0 = \{(x, \Phi(x)) | x \in [x_0 - \varepsilon, x_0 + \varepsilon]\}$ and $\Gamma_0 = \{(\xi, z) \in \mathbb{R}^2 \times \mathbb{R} |\, 1 \leq z \leq 2,\ \frac{\xi}{z} \in \gamma_0\}$, and correspondingly $\tilde{\gamma}_0 = \{(x, \tilde{\Phi}(x)) | x \in [\varepsilon, \varepsilon]\}$ and $\tilde{\Gamma}_0$, then we compute

$$\tilde{\Gamma}_0 = A\Gamma_0$$

with the volume preserving transformation

$$A = \begin{pmatrix} 1 & 0 & -x_0 \\ -\Phi'(x_0) & 1 & x_0\Phi'(x_0) - \Phi(x_0) \\ 0 & 0 & 1 \end{pmatrix}.$$

This justifies (6).
Additionally, we split $\Phi|_{(-\varepsilon,\varepsilon)}$ into $\Phi|_{(-\varepsilon,0)}$ and $\Phi|_{(0,\varepsilon)}$. On $(-\varepsilon, 0)$, we switch to the function $\Phi^- : (0, \varepsilon) \to \mathbb{R}, \Phi^-(x) = \Phi(-x)$, and hence, we need just to consider parts like $\Phi|_{(0,\varepsilon)}$. Eventually, without loss of generality, we may assume that $\Phi^{(m)}(0) > 0$. To summarize, we have

$$\Phi^{(m)}(0) > 0 \qquad \text{and} \qquad \Phi(0) = \Phi'(0) = \ldots = \Phi^{(m-1)}(0) = 0. \tag{7}$$

By choosing ε small enough we may ensure that $\Phi^{(m)}|_{[0,\varepsilon]} > 0$. The next step is to establish the following:

$$\forall k \leq m\ \forall 0 < x \leq \varepsilon :\ \Phi^{(k)}(x) > 0. \tag{8}$$

If this doesn't hold, we would find $x_k > 0$ such that $\Phi^{(k)}(x_k) \leq 0$. Since $\Phi^{(k)}(0) = 0$, the mean value theorem would provide a $x_{k+1} \in (0, x_k)$ such that $\Phi^{(k+1)}(x_{k+1}) \leq 0$, and by induction eventually a $x_m > 0$ such that $\Phi^{(m)}(x_m) \leq 0$ in contradiction to our assumption.
Furthermore $\Phi(x) \approx x^m$. To be more precise, Taylor expansion provides

$$\frac{1}{m!} \min_{t \in [0,\varepsilon]} \Phi^{(m)}(t)\ x^m \leq \Phi(x) \leq \frac{1}{m!} \max_{t \in [0,\varepsilon]} \Phi^{(m)}(t)\ x^m,$$

where the minimum on the left-hand-side doesn't vanish, since (7) and (8) imply $\Phi^{(m)}(t) > 0$ for all $t \in [0, \varepsilon]$. In other words, we find a $\chi \in C^2([0, \varepsilon], (0, \infty))$ fulfilling $\Phi(x) = x^m\chi(x)$. This implies

$$\Phi'(x) = mx^{m-1}\chi(x) + x^m\chi'(x) = x^{m-1}[m\chi(x) + x\chi'(x)].$$

By reducing ε if necessary, we may assume $\chi^1 := m\chi(x) + x\chi'(x) \in C^1([0, \varepsilon], (0, \infty))$. In the same manner, we find $\chi^2 \in C([0, \varepsilon], (0, \infty))$ fulfilling

$$\Phi''(x) = x^{m-2}\chi^2(x).$$

By further reduction of ε, we may also find an open neighbourhood of the origin, were Φ has the desired properties. Eventually, we may rescale $\varepsilon = 1$.

2 Reduction of the problem

Altogether, it remains to discuss the following problem:

Definition
Let $m \in \{3, 4, \ldots\}$ *and let* $\Phi : [0,1] \to \mathbb{R}$ *satisfiy the following conditions:*

(i) $\exists \chi \in C^2[0,1], \chi > 0 : \Phi(x) = x^m \chi(x)$,
$\exists \chi^1 \in C^1[0,1], \chi^1 > 0 : \Phi'(x) = x^{m-1}\chi^1(x)$,
$\exists \chi^2 \in C[0,1], \chi^2 > 0 : \Phi''(x) = x^{m-2}\chi^2(x)$,

(ii) $\Phi^{(k)}(x) > 0$ *für* $x > 0$, $k \leq m$*; especially,* Φ *and* Φ' *are convex.*

We define the generalised cone (more exactly, a section of a cone)

$$\Gamma = \{(x, y, z) \in \mathbb{R}^3 |\, 0 \leq x \leq 1,\ 1 \leq z \leq 2,\ \Phi\left(\frac{x}{z}\right) = \frac{y}{z}\}.$$

The associated surface measure will be denoted by σ*.*

Theorem 2.1
For $1 \leq p < \frac{m+1}{m}$ *and* $\frac{1}{q} \geq \frac{m+1}{p'}$ *holds*

$$\|\hat{f}|_\Gamma\|_{L_q(\Gamma,\sigma)} \leq C\|f\|_{L_p(\mathbb{R}^3)} \qquad \forall f \in \mathcal{S}(\mathbb{R}^3). \tag{9}$$

2.1 Decomposition

The critical part of Γ is the origin, where the curvature vanishes. To accommodate this fact, we decompose Γ in dyadic pieces, becoming smaller near the origin. By rescaling to the case of (almost) constant curvature, we would be able to make use of already known estimates. Unfortunately, summation is only possible if $\frac{1}{q} > \frac{m+1}{p'}$. In the limit case $\frac{1}{q} = \frac{m+1}{p'}$, we need to deal with the problem as a whole. For this, we will further decompose each dyadic piece, depending on the curvature.
Let $\Gamma^\delta = \{(x, y, z) \in \mathbb{R}^3 |\, 0 \leq x < 1,\ 1 \leq z \leq 2,\ \Phi\left(\frac{x}{z}\right) \leq \frac{y}{z} \leq \Phi\left(\frac{x}{z}\right) + \delta\}$ be the thickening of Γ by $\delta > 0$ (we drop the points with $x = 1$ for technical purposes). Moving on the $x-$axis from the origin by length $\sqrt[m]{\delta}$ corresponds to Φ changing by δ. In other words, this part of Γ is contained in a box of width δ. We thus define $\gamma = \sqrt[m]{\delta}$ and

$$\Gamma^\delta_k = \{(x, y, z) \in \Gamma^\delta | (2^k\text{-}1)\gamma \leq x < (2^{k+1}\text{-}1)\gamma\}, \qquad k = 0, \ldots, \left\lceil \log_2 \frac{1}{\gamma} \right\rceil - 1.$$

Now how to determine the finer decomposition? We change coordinates, or respectively Φ by affine transformation into

$$\Phi^k(x) = \Phi(x + (2^k\text{-}1)\gamma) - \Phi((2^k\text{-}1)\gamma) - x\Phi'((2^k\text{-}1)\gamma),$$

such that

$$\Phi^k(0) = 0 = (\Phi^k)'(0).$$

According to Taylor, we get

$$\Phi^k(x) \approx \frac{m(m-1)}{2}((2^k\text{-}1)\gamma)^{m-2}x^2 + \mathcal{O}(x^3).$$

On which distance γ_k from the (new) origin does Φ varys at most δ? We demand ($k \neq 0$)

$$\begin{aligned} \delta =& \Phi^k(\gamma_k) \approx \frac{m(m-1)}{2}((2^k\text{-}1)\gamma)^{m-2}\gamma_k^2, \\ \text{i.e.} \qquad \gamma_k \approx& \sqrt{\delta((2^k\text{-}1)\gamma)^{2-m}} \approx 2^{k(1-\frac{m}{2})}\gamma. \end{aligned}$$

Concerning the height, we decompose equidistantly with width β, where we require $\beta \lesssim \delta$. This ensures that the projection of such a set in x-$y-$space does not appear to different to a intersection parallel to x-$y-$space. To choose $\beta = \delta$ would be appropriate and you might assume this. Nevertheless, we distinct these two quantities to be aware how each of them effects our computations. We will see that the β's cancels against each others at the end, reflecting the fact that there is no impact from the z-dircetion. We obtain a decomposition of Γ^δ as follows:

DEFINITION (DECOMPOSITION)
Let $\delta > 0$, $\gamma > 0$ and $\gamma^m = \delta$, let $\beta < \delta$ with $\frac{1}{\beta} \in \mathbb{N}$. For $k = 0, \ldots, |\log_2 \gamma| - 1$, $j \in I_k = \{0, \ldots, 2^{k\frac{m}{2}} - 1\}$ and $n = \frac{1}{\beta}, \ldots \frac{2}{\beta} - 1$ define $\gamma_k = 2^{k(1-\frac{m}{2})}\gamma$, $x_{kj} = (2^k\text{-}1)\gamma + j\gamma_k$, $x_k = x_{k,0}$ and

$$\Gamma^\delta_{kjn} = \{(x,y,z) \in \mathbb{R}^3 |\, n\beta \leq z \leq (n+1)\beta,\ \Phi\left(\tfrac{x}{z}\right) \leq \tfrac{y}{z} \leq \Phi\left(\tfrac{x}{z}\right) + \delta,\ x_{kj} \leq x < x_{k,j+1}\}. \tag{10}$$

Furthermore let ϕ_{kjn} be a bump function adapted to Γ_{kjn}. To be more precise, if $\eta \in C_0^\infty(\mathbb{R})$, $\chi_{[-1,1]} \leq \eta \leq \chi_{[-\frac{5}{4},\frac{5}{4}]}$, let

$$\phi_{kjn}(x,y,z) = \eta\left(\frac{x - x_{kj}}{\gamma_k}\right)\eta\left(\frac{y - z\Phi(x/z)}{\delta}\right)\eta\left(\frac{z - n\beta}{\beta}\right).$$

This means that ϕ_{kjn} is to some extend supported in an anisotropic thickening of Γ^δ_{kjn}, precisely in the set

$$\begin{aligned} \tilde{\Gamma}^\delta_{kjn} =& \{(x,y,z) \in \mathbb{R}^3 |\, |z - n\beta| \leq \tfrac{5}{4}\beta,\ |\tfrac{y}{z} - \Phi\left(\tfrac{x}{z}\right)| \leq \tfrac{5}{4}\delta,\ |x - x_{kj}| \leq \tfrac{5}{4}\gamma_k\} \\ \subset& \bigcup_{u,v,w \in \{-1,0,1\}} (u\gamma_k, v\delta, w\beta) + \Gamma^\delta_{kjn}. \end{aligned}$$

For the further proceeding, we will always denote by α the tripel (k,j,n), and, if required, by μ a second tripel (l,i,p).

For simplification, we write $\sum_\alpha$ instead of $\sum\limits_{k=0}^{|\log_2\gamma|-1} \sum\limits_{j=0}^{2^{k\frac{m}{2}}-1} \sum\limits_{n=\frac{1}{\beta}}^{\frac{2}{\beta}-1}$.



Fig. 1: Decomposition

2.2 Proceeding to the crux of the problem

Of essential impact is the following theorem. It is a weighted, discrete version of the adjoint restriction estimate and, as we will see, already implies the restriction theorem.

Theorem 2.2
Let $p' > m+1$, $\frac{3}{p'} < \frac{1}{q} < \frac{1}{2} + \frac{1}{p'}$. *Then*

$$\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'} \lesssim \delta^{\frac{1}{q}} \| \sum_{\alpha=(k,j,n)} a_\alpha ((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}} \phi_\alpha \|_{q'}. \tag{11}$$



Fig. 2: Range of p' and q

The proof will be the main work in this paper and be done in the next chapters. First of all, we will derive the retriction theorem.
As an immediate consequence of theorem 2.2, we get as corollary

Corollary 2.3
Let $p' > m+1$, $\frac{3}{p'} < \frac{1}{q}$. *Then*

$$\|\sum_{\alpha} a_\alpha \hat{\phi}_\alpha\|_{p'} \lesssim \delta^{\frac{1}{q}} \|\sum_{\alpha=(k,j,n)} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}}\phi_\alpha\|_{q',1}. \tag{11}$$

Proof: Actually, we just need to use that (11) is valid for every p as above and for some range $\frac{3}{p'} < \frac{1}{q} < \frac{3}{p'} + \varepsilon_p$. According to the assumptions, we have $p' > m+1 \geq 4$. Hence $\frac{3}{p'} = \frac{2}{p'} + \frac{1}{p'} < \frac{1}{2} + \frac{1}{p'}$. Therefore we find r with $\frac{3}{p'} < \frac{1}{r} < \frac{1}{2} + \frac{1}{p'}$ (cf. figure 2), and thus the requirements of theorem 2.2. So it is sufficient to show that if (11) holds for some (p,r), then it also holds for (p,q) with $\frac{1}{r} < \frac{1}{q}$. Under this condition, there exists $1 \leq s < \infty$ such that $\frac{1}{r'} = \frac{1}{q'} + \frac{1}{s}$. This means

$$\frac{1}{r} - \frac{1}{q} = \frac{1}{q'} - \frac{1}{r'} = -\frac{1}{s}. \tag{12}$$

For (x,y,z) in the support of ϕ_{kjn}, it follows $x \approx (2^k\text{-}1)\gamma$, so we introduce $g(x,y,z) := x^{-\frac{1}{s}}$. We now may apply theorem 2.2 and use Hölders inequality for Lorentz spaces (see Lemma 7.2):

$$\begin{aligned}
\|\sum_{\alpha} a_\alpha \hat{\phi}_\alpha\|_{p'} \lesssim& \delta^{\frac{1}{r}} \|\sum_{\alpha=(k,j,n)} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{r}-\frac{m+1}{p'}}\phi_\alpha\|_{r'} \\
=& \delta^{\frac{1}{r}} \|\sum_{\alpha=(k,j,n)} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}-\frac{1}{s}}\phi_\alpha\|_{r'} \\
\lesssim& \delta^{\frac{1}{r}} \|\sum_{\alpha} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}}\phi_\alpha g\|_{L_{r'}(\Gamma^\delta)} \\
\leq& \delta^{\frac{1}{r}} \|\sum_{\alpha} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}}\phi_\alpha\|_{L_{q',r'}(\Gamma^\delta)} \|g\|_{L_{s,\infty}(\Gamma^\delta)}.
\end{aligned} \tag{13}$$

A short computation yields

$$\begin{aligned}
|\{(x,y,z) \in \Gamma^\delta : x^{-\frac{1}{s}} > \lambda\}| =& \int_0^{\lambda^{-s}} \int_1^2 \int_{z\Phi(\frac{x}{z})}^{z\Phi(\frac{x}{z})+z\delta} \mathrm{d}y\ \mathrm{d}z\ \mathrm{d}x \\
=& \delta\lambda^{-s} \int_1^2 z\ \mathrm{d}z \approx \delta\lambda^{-s},
\end{aligned}$$

i.e.

$$\|g\|_{L_{s,\infty}(\Gamma^\delta)} = \sup_{\lambda>0} \lambda |\{(x,y,z) \in \Gamma^\delta : x^{-\frac{1}{s}} > \lambda\}|^{\frac{1}{s}} \approx \delta^{\frac{1}{s}}. \tag{14}$$

Combined with (13), we end up with

$$\begin{aligned}
\|\sum_{\alpha} a_\alpha \hat{\phi}_\alpha\|_{p'} \lesssim& \delta^{\frac{1}{r}+\frac{1}{s}} \|\sum_{\alpha} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}}\phi_\alpha\|_{q',r'} \\
\leq& \delta^{\frac{1}{q}} \|\sum_{\alpha} a_\alpha((2^k\text{-}1)\gamma)^{\frac{1}{q}-\frac{m+1}{p'}}\phi_\alpha\|_{q',1}\ ,
\end{aligned}$$

since (12) means $\frac{1}{r} + \frac{1}{s} = \frac{1}{q}$. □

For $\frac{1}{q} = \frac{m+1}{p'}$ as considered in theorem 2.1, (11) reads

$$\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'} \lesssim \delta^{\frac{1}{q}} \| \sum_{\alpha} a_\alpha \phi_\alpha \|_{(q',1)} \tag{15}$$

for all sequences a_α and every $\delta > 0$. In the next chapter, we will conclude that this statement, at least when $q < \infty$, implies the desired restriction theorem. To be more precise, for $\frac{1}{q} = \frac{m+1}{p'}$ and $q < \infty$ or $p' > m+1$ respectively, the weak $L_p(\mathbb{R}^3)$-$L_{q,\infty}(\Gamma)$ estimate

$$\| \hat{f}|_\Gamma \|_{L_{q,\infty}(\Gamma)} \lesssim \| f \|_p \qquad \forall f \in \mathcal{S}(\mathbb{R}^3)$$

holds. Eventually we finish the proof by using Marcinkiewicz interpolation. In the endpoint $p = 1$, $q = \infty$, where we are not able to interpolate, the estimate is trivial.

3 Dense families

3.1 Basic properties

Definition (Dense family)
Let X be a metric (or even topological) space. A family M_δ, $\delta > 0$, of subsets of X is called a dense family in X, *if*

$$\forall\ f \in X\ \exists \delta_f > 0\ \forall 0 < \delta < \delta_f\ \exists f_\delta \in M_\delta\ f = \lim_{\delta\to 0} f_\delta.$$

Remark
Given a dense family M_δ, $\delta > 0$, the set $\bigcup\limits_{\delta>0} M_\delta$ is dense in X. Conversely, none of the sets M_δ has to be dense.

Definition
Let (X, d) be a metric space.

- *For a given set A with $diam(A) < \infty$, we define its doubling by $\tilde{A} = \bigcup\limits_{x\in A} B(x, diam\, A)$. Observe that $diam\, \tilde{A} \le 3\ \ diam\, A$.*
- *A family $\{A_i\}_i$ of subsets of X is said to fulfill the* bounded overlap property *if for some constant $N \in \mathbb{N}$, every point of X is contained in at most N sets A_i.*

We will discuss an example of a dense family:

Lemma 3.1
Let X be a compact metric space and $\delta_0 > 0$. For every $\delta \in (0, \delta_0)$, let $\{A_{\delta,i}\}_i$ be a family with the bounded overlap property *covering X, and satisfying*

$$\lim_{\delta\to 0} \sup_i diam(A_{\delta,i}) = \ \ 0. \tag{16}$$

Let $\{\phi_{\delta,i}\}_i$ be a partition of unity adopted to $\{A_{\delta,i}\}_i$, i.e.

$$\sum_i \phi_{\delta,i} = 1 \text{ on } X, \qquad \phi_{\delta,i} \in C(X, [0,\infty)) \qquad \text{and} \qquad supp\, \phi_{\delta,i} \subset \tilde{A}_{\delta,i}.$$

Under these assumptions, $M_\delta := LH\{\phi_{\delta,i}\}_i$, $\delta > 0$ is a dense family in $(C(X), \|\cdot\|_\infty)$.

Proof: Let $f \in C(X)$, $\delta_f := \delta_0$ and $x_{\delta,i} \in A_{\delta,i}$. Define

$$f_\delta := \sum_i f(x_{\delta,i})\phi_{\delta,i} \in M_\delta.$$

We have to show that $\lim\limits_{\delta\to 0} \|f_\delta - f\|_\infty = 0$, or

$$\forall \varepsilon > 0\ \exists \delta(\varepsilon) > 0\ \forall \delta \in (0, \delta(\varepsilon))\ \|f_\delta - f\|_\infty < \varepsilon$$

respectively. The compactness of X provides uniformly continuity of f. Given $\varepsilon > 0$ there exists $\gamma > 0$, such that $|f(x) - f(y)| < \varepsilon$ for all $x, y \in X$ with distance $d(x, y) < \gamma$. Using condition (16), we find $\delta(\varepsilon) > 0$ such that $\sup_i diam(A_{\delta,i}) < \frac{1}{3}\gamma$ for all $0 < \delta < \delta(\varepsilon)$, i.e. $\forall i: \ diam(\tilde{A}_{\delta,i}) < \gamma$, for all $\delta \in (0, \delta(\varepsilon))$. Therefore, we obtain

$$\forall \delta \in (0, \delta(\varepsilon)) \ \forall i \ \forall x, y \in \tilde{A}_{\delta,i} \ |f(x) - f(y)| < \varepsilon. \tag{17}$$

Since $supp\, \phi_{\delta,i} \subset \tilde{A}_{\delta,i}$, we conclude

$$\forall \delta \in (0, \delta(\varepsilon)) \ \forall i \ \forall y \in \tilde{A}_{\delta,i} \forall x \in X \ |f(x) - f(y)|\phi_{\delta,i}(x) \leq \varepsilon \phi_{\delta,i}(x). \tag{18}$$

For such a δ and every $x \in X$

$$\begin{aligned} |f_\delta(x) - f(x)| &= \left| \sum_i f(x_{\delta,i})\phi_{\delta,i}(x) - f(x) \sum_i \phi_{\delta,i}(x) \right| \\ &\leq \sum_i |f(x_{\delta,i}) - f(x)|\phi_{\delta,i}(x) \\ &\underset{(18)}{<} \varepsilon \sum_i \phi_{\delta,i}(x) = \varepsilon \end{aligned}$$

holds, as we claimed. □

EXAMPLE

We consider as metric space the surface

$$X = \Gamma = \left\{ (x, y, z) \in \mathbb{R}^3 | \, 0 \leq x \leq 1, \ 1 \leq z \leq 2, \ \Phi\left(\frac{x}{z}\right) = \frac{y}{z} \right\}$$

from the previous chapter. The families

$$A_{\delta,kjn} = \Gamma \cap \Gamma^\delta_{kjn} = \{(x, y, z) \in \mathbb{R}^3 | \, n\beta \leq z \leq (n+1)\beta, \ \Phi\left(\tfrac{x}{z}\right) = \tfrac{y}{z}, \ x_{kj} \leq x < x_{k,j+1}\},$$

$x_{kj} = (2^k\text{-}1)\gamma + j2^{k(1-\frac{m}{2})}\gamma$, *are fulfilling the required assumptions of Lemma 3.1: For every* δ, *they present a covering of* X *(with their overlap having measure zero) and it holds*

$$\begin{aligned} \sup_{kjn} diam(A_{\delta,kjn}) &= \sup_{kjn} \max\{\beta, 2^{k(1-\frac{m}{2})}\gamma\} \\ &= \max\{\beta, \gamma\} \leq \max\{\delta, \delta^{\frac{1}{m}}\} \\ &= \delta^{\frac{1}{m}} \overset{\delta \to 0}{\longrightarrow} 0. \end{aligned}$$

The sets Γ^δ_{kjn} *can be seen as a certain thickening of* $A_{\delta,kjn}$ *by* δ *in the direction of* $(0, 1, 0)$, *which we are going to formalise soon.*

3.2 An application

A classical result states that it is enough to know the behaviour of a bounded operator on some dense subset. We will obtain a similar statement concerning dense families.
During the following explanations, let Γ be a compact hypersurface in $\mathbb{R}^n$ and σ the associated surface measure. Moreover, we take the existence of a universal transversal unit vector $N \in \mathbb{R}^n$ for granted, i.e. $\forall p \in \Gamma: \ N \notin T_p\Gamma$. (This is not a strong assumption, since Γ can at least be decomposed in finitely many subsets with such a universal transversal vector.)

Definition (δ-thickening)

Let Γ^δ be the thickening of Γ by $\delta > 0$ in the direction of the universal transversal vector N, i.e. $\Gamma^\delta = \Gamma + [0,\delta]N$.
Given $h \in L_\infty(\Gamma)$, we denote by $\tilde{h}$ the continuation along N on Γ^δ, defined by $\tilde{h}(\xi + tN) = h(\xi)\eta\left(\frac{t}{\delta}\right)$ for all $\xi \in \Gamma$, $t \in \mathbb{R}$. Here $\eta \in C_0^\infty(\mathbb{R})$ mit $\chi_{[-1,1]} \leq \eta \leq \chi_{[-2,2]}$.

Example

For our example, $N = (0,1,0)$ is a universal transversal unit vector. The functions

$$\phi_{kjn}(x,y,z) = \eta\left(\frac{x - x_{kj}}{\gamma_k}\right)\eta\left(\frac{y - z\Phi(x/z)}{\delta}\right)\eta\left(\frac{z - n\beta}{\beta}\right)$$

introduced in chapter 2.1 can be considered as the continuation of the functions

$$\phi_{kjn}(x, z\Phi(x/z), z) = \eta\left(\frac{x - x_{kj}}{\gamma_k}\right)\eta\left(\frac{z - n\beta}{\beta}\right),$$

which are defined just on Γ.

Remark 3.2

Integration on Γ^δ can be done as follows:
There is a constant $A > 0$ and a smooth density $\psi : \Gamma \to (\frac{1}{A}, A)$ in the way that

$$\int_{\Gamma^\delta} f(x)\ \mathrm{d}x = \int_0^\delta \int_\Gamma f(\xi + tN)\psi(\xi)\ \mathrm{d}\sigma(\xi)\ \mathrm{d}t \qquad \forall f \in L_1(\Gamma^\delta).$$

Introducing $\mathrm{d}\mu = \psi d\sigma$, the formula gives

$$\int_{\Gamma^\delta} f(x)\ \mathrm{d}x = \int_0^\delta \int_\Gamma f(\xi + tN)\ \mathrm{d}\mu(\xi)\ \mathrm{d}t \qquad \forall f \in L_1(\Gamma^\delta). \tag{19}$$

For our purposes, it's no matter wether to consider σ or μ: If the conjugated retriction theorem

$$\|\widehat{g d\mu}\|_{L_p(\mathbb{R}^n)} \leq C\|g\|_{L_q(\Gamma)} \qquad \forall g \in C(\Gamma),$$

holds, then

$$\|\widehat{g d\sigma}\|_{L_p(\mathbb{R}^n)} = \left\|\widehat{\frac{g}{\psi} d\mu}\right\|_{L_p(\mathbb{R}^n)} \leq C\left\|\frac{g}{\psi}\right\|_{L_q(\Gamma)} \leq AC\ \|g\|_{L_q(\Gamma)} \qquad \forall g \in C(\Gamma)$$

holds as well.

Lemma 3.3

Let $C > 0$, $\delta_0 > 0$ and $1 \leq p \leq \infty$, $1 < q \leq \infty$.
Furthermore let $\{M_\delta\}_{\delta > 0}$ be a dense family in $C(\Gamma)$ fulfilling

$$\left\|\hat{\tilde{h}}\right\|_{L_{p'}(\mathbb{R}^n)} \leq C\delta^{\frac{1}{q}}\left\|\tilde{h}\right\|_{L_{q',1}(\Gamma^\delta)} \qquad \forall\delta \in (0,\delta_0)\ \forall h \in M_\delta. \tag{20}$$

Then

$$\|\widehat{g d\sigma}\|_{L_{p'}(\mathbb{R}^n)} \leq C'\|g\|_{L_{q',1}(\Gamma,\sigma)} \qquad \forall g \in C(\Gamma)$$

and

$$\|\hat{f}|_\Gamma\|_{L_{q,\infty}(\Gamma,\sigma)} \leq C'\|f\|_{L_p(\mathbb{R}^n)} \qquad \forall f \in \mathcal{S}(\mathbb{R}^n) \subset L_p(\mathbb{R}^n). \tag{21}$$

A more canonical version of the lemma would be to test the estimate on a *dense set* like $\bigcup\limits_{\eta>0} M_\eta$. To handle the surface measure, we then may use a δ-thickening of Γ. Thus the canonical version of condition (20) would be

$$\left\|\hat{\tilde{h}}\right\|_{L_{p'}(\mathbb{R}^n)} \leq C\delta^{\frac{1}{q'}}\left\|\tilde{h}\right\|_{L_{q',1}(\Gamma^\delta)} \qquad \forall\delta\in(0,\delta_0)\ \forall h\in\bigcup_{\eta>0}M_\eta .$$

The lemma states that it suffies to check the estimate for less functions: We are allowed to link up η and δ. These considerations eventually lead to the definition of a dense family.

Proof:
According to remark 3.2, we may use the measure μ instead of σ.
By our assumptions, every $g\in C(\Gamma)$ provides approximations $g_\delta\in M_\delta$, $\delta>0$ such that $\lim\limits_{\delta\to 0} g_\delta = g$ in $L_\infty(\Gamma)$.
First of all, we observe $\lim\limits_{\delta\to 0}\widehat{g_\delta \mathrm{d}\mu} = \widehat{g\mathrm{d}\mu}$ uniformly and pointwise, because

$$\left\|\widehat{g_\delta\mathrm{d}\mu}-\widehat{g\mathrm{d}\mu}\right\|_\infty = \sup_{x\in\mathbb{R}^n}\Big|\int_\Gamma (g_\delta-g)(\xi)e^{-ix\xi}\ \mathrm{d}\mu(\xi)\Big| \leq \mu(\Gamma)\|g_\delta-g\|_\infty \overset{\delta\to 0}{\longrightarrow} 0.$$

This implies

$$\begin{aligned}
\frac{1}{\delta}\hat{\tilde{g}}_\delta(x) &\overset{(19)}{=}\frac{1}{\delta}\int_{-\infty}^{\infty}\int_\Gamma \tilde{g}_\delta(\xi+tN)e^{-ix\cdot(\xi+tN)}\ \mathrm{d}\mu(\xi)\ \mathrm{d}t\\
&=\int_\Gamma g_\delta(\xi)e^{-ix\cdot\xi}\ \mathrm{d}\mu(\xi)\quad \frac{1}{\delta}\int_{-\infty}^{\infty}\chi\left(\frac{t}{\delta}\right)e^{-itx\cdot N}\ \mathrm{d}t\\
&=\underbrace{\widehat{g_\delta\mathrm{d}\mu(x)}}_{\overset{\delta\to 0}{\longrightarrow}\widehat{g\mathrm{d}\mu}(x)}\quad\underbrace{\frac{1}{\delta}\int_{-\infty}^{\infty}\chi\left(\frac{t}{\delta}\right)e^{-itx\cdot N}\ \mathrm{d}t}_{\overset{\delta\to 0}{\longrightarrow}\|\chi\|_1}\qquad\overset{\delta\to 0}{\longrightarrow}\quad \|\chi\|_1\widehat{g\mathrm{d}\mu}(x),
\end{aligned}$$

i.e.

$$\widehat{g\mathrm{d}\mu}(x)\approx\lim_{\delta\to 0}\frac{1}{\delta}\hat{\tilde{g}}_\delta(x). \tag{22}$$

Furthermore, we compute the distribution function

$$\begin{aligned}
d_{\tilde{g}_\delta}(s) =&\left|\{v\in\Gamma^\delta:\ |\tilde{g}_\delta(v)|>s\}\right|\\
\overset{(19)}{=}&(\mu\otimes\lambda)\Big(\{(\xi,t)\in\Gamma\times\mathbb{R}:\ |\underbrace{\tilde{g}_\delta(\xi+tN)}_{=g_\delta(\xi)\chi\left(\frac{t}{\delta}\right)}|>s\}\Big)\\
\leq&4\delta\mu\left(\{(\xi\in\Gamma:\ |g_\delta(\xi)|>s\}\right)=4\delta d_{g_\delta}(s)
\end{aligned}$$

and thus the decreasing rearrangement

$$\tilde{g}_\delta^*(t)=\inf\{s>0|d_{\tilde{g}_\delta}(s)\leq t\}\leq\inf\left\{s>0|d_{g_\delta}(s)\leq\frac{t}{4\delta}\right\}=g_\delta^*\left(\frac{t}{4\delta}\right).$$

These estimates yield

$$\begin{aligned}
\|\tilde{g}_\delta\|_{L_{q',1}(\Gamma^\delta)} =& \int_0^\infty t^{\frac{1}{q'}} \tilde{g}_\delta^*(t) \frac{\mathrm{d}t}{t} \\
\leq& \int_0^\infty t^{\frac{1}{q'}} g_\delta^* \left(\frac{t}{4\delta}\right) \frac{\mathrm{d}t}{t} \\
\approx& \delta^{\frac{1}{q'}} \int_0^\infty t^{\frac{1}{q'}} g_\delta^*(t) \frac{\mathrm{d}t}{t} \\
=& \delta^{\frac{1}{q'}} \big\|g_\delta\big\|_{L_{q',1}(\Gamma,\mu)}. \qquad (23)
\end{aligned}$$

Moreover, it is valid for a general $f \in L_\infty(\Gamma)$ that

$$\begin{aligned}
\|f\|_{L_{q',1}(\Gamma,\mu)} =& \int_0^\infty t^{\frac{1}{q'}} f^*(t) \frac{\mathrm{d}t}{t} \\
=& \int_0^{\mu(\Gamma)} t^{\frac{1}{q'}-1} f^*(t) \ \mathrm{d}t \\
\leq& f^*(0) \int_0^{\mu(\Gamma)} t^{\frac{1}{q'}-1} \ \mathrm{d}t \\
\approx& \|f\|_\infty,
\end{aligned}$$

since $\frac{1}{q'} > 0$, i.e. $q > 1$, and therefore

$$\|g_\delta - g\|_{L_{q',1}(\Gamma,\mu)} \lesssim \|g_\delta - g\|_\infty \overset{\delta\to 0}{\longrightarrow} 0,$$

so especially

$$\lim_{\delta\to 0} \|g_\delta\|_{L_{q',1}(\Gamma,\mu)} = \|g\|_{L_{q',1}(\Gamma,\mu)}. \qquad (24)$$

Hence by using assumption (20), we obtain

$$\begin{aligned}
\big\|\widehat{g\mathrm{d}\mu}\big\|_{L_{p'}(\mathbb{R}^n)} \overset{(22)}{\approx}& \ \big\|\liminf_{\delta\to 0} \frac{1}{\delta}\widehat{\tilde{g}_\delta}\big\|_{L_{p'}(\mathbb{R}^n)} \\
\overset{Fatou}{\leq}& \ \liminf_{\delta\to 0} \delta^{-1}\big\|\widehat{\tilde{g}_\delta}\big\|_{L_{p'}(\mathbb{R}^n)} \\
\overset{(20)}{\leq}& \ C\liminf_{\delta\to 0} \delta^{\frac{1}{q}-1}\big\|\tilde{g}_\delta\big\|_{L_{q',1}(\Gamma^\delta)} \\
\overset{(23)}{\lesssim}& \ C\liminf_{\delta\to 0} \delta^{\frac{1}{q}-1+\frac{1}{q'}}\big\|g_\delta\big\|_{L_{q',1}(\Gamma,\mu)} \\
\overset{(24)}{=}& \ C\big\|g\big\|_{L_{q',1}(\Gamma,\mu)}.
\end{aligned}$$

The second part of our claim is a simple duality argument. □

4 Estimation of the overlap

4.1 Straightforward results

It will become essential for us to understand the overlap of the sets $\Gamma^\delta_{kjn} + \Gamma^\delta_{lip}$, $k, l = 0, \ldots, |\log_2 \gamma| - 1$, $j \in I_k$, $i \in I_l$, $n, p = \frac{1}{\beta}, \ldots, \frac{2}{\beta} - 1$, i.e. to examine the maximal number of them containing a single point. However, this number will not be bounded by a absolute constante (which would be optimal). We will get some overlap from the $z-$direction. Nevertheless, this will not cause us any trouble.

Lemma 4.1
For every $\xi \in \mathbb{R}^3$ holds

$$\#\{(n,p) | \exists k, l, j, i : \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} \leq 3\beta^{-1}. \tag{25}$$

Proof: If $\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu$, there exist $\mathbf{x}_1 \in \Gamma^\delta_{kjn}$ and $\mathbf{x}_2 \in \Gamma^\delta_{lip}$ with $\xi = \mathbf{x}_1 + \mathbf{x}_2$. Let z_1, z_2 denote the last component of $\mathbf{x}_1$ and $\mathbf{x}_2$ respectively, which means $n\beta \leq z_1 \leq (n+1)\beta$ and $p\beta \leq z_2 \leq (p+1)\beta$. The sum of both inequalities leads to $(n+p)\beta \leq z \leq (n+p+2)\beta$. This implies $\frac{z}{\beta} - 2 \leq n + p \leq \frac{z}{\beta}$ and hence

$$n \in -p + [\tfrac{z}{\beta} - 2, \tfrac{z}{\beta}] \cap \mathbb{N}.$$

We conclude

$$\begin{aligned}
\#\{(n,p) | \exists k, l, j, i : \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} &= \sum_{p=\frac{1}{\beta}}^{\frac{2}{\beta}-1} \#\{n | \exists k, l, j, i : \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} \\
&\leq \sum_{p=\frac{1}{\beta}}^{\frac{2}{\beta}-1} \#\left([\tfrac{z}{\beta} - 2, \tfrac{z}{\beta}] \cap \mathbb{N}\right) \\
&\leq \sum_{p=\frac{1}{\beta}}^{\frac{2}{\beta}-1} 3 = 3\beta^{-1},
\end{aligned}$$

completing the proof. □

Exploiting the dyadic structur of the decomposition, we obtain a further simple result:

Lemma 4.2
For $k = 0, \ldots, |\log_2 \gamma| - 1$ let $V_k(n,p) = \bigcup\{\Gamma^\delta_{kjn} + \Gamma^\delta_{lip} | l \leq k,\ j \in I_k,\ i \in I_l\}$. Then for all $\xi \in \mathbb{R}^3$, $n, p = \frac{1}{\beta}, \ldots, \frac{2}{\beta} - 1$ holds

$$\#\{k | \xi \in V_k(n,p)\} \leq 3.$$

Proof: The procedure is comparable to the previous proof, though we now concentrate on the x-component. If $\xi \in V_k(n,p) = \bigcup\{\Gamma^\delta_{kjn} + \Gamma^\delta_{lip} | l \leq k,\ j \in I_k,\ i \in I_l\}$ and x denotes the first component of ξ, there exist l, j, i such that

$$(2^k\text{-}1)\gamma + j\gamma_k + (2^l\text{-}1)\gamma + i\gamma_l \leq x \leq (2^k\text{-}1)\gamma + (j+1)\gamma_k + (2^l\text{-}1)\gamma + (i+1)\gamma_l.$$

Since we are just interested in the dyadic size, we estimate quite roughly

$$(2^k\text{-}1)\gamma \leq x \leq (2^k\text{-}1)\gamma + 2^{k\frac{m}{2}} 2^{k(1-\frac{m}{2})}\gamma + (2^l\text{-}1)\gamma + 2^{l\frac{m}{2}} 2^{l(1-\frac{m}{2})}\gamma = (2^{k+1}\text{-}1)\gamma + (2^{l+1}\text{-}1)\gamma.$$

and since we assumed $l \leq k$, this reduces to

$$(2^k\text{-}1)\gamma \leq x \leq (2^{k+1}\text{-}1)\gamma + (2^{l+1}\text{-}1)\gamma \leq (2^{k+2}\text{-}1)\gamma.$$

It follows

$$2^k \leq \frac{x}{\gamma} + 1 \leq 2^{k+2}$$

and therefore

$$\log_2\left(\frac{x}{\gamma} + 1\right) - 2 \leq k \leq \log_2\left(\frac{x}{\gamma} + 1\right).$$

Our claim is an immediate consequence of this inequality. □

The handling of the overlap concerning the remaining parameters is more complicated. Here we need to involve the $y-$coordinate. Therefore, we need as a start a new coordinate system.

4.2 Further results

Lemma 4.3

There exists an absolute constant $C > 0$ (not depending on δ) such that for all $\xi \in \mathbb{R}^3$, $k = 0, \ldots, |\log_2 \gamma| - 1$, $n, p = \frac{1}{\beta}, \ldots, \frac{2}{\beta} - 1$

$$\#\{(j,l,i) | \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} \leq C. \tag{26}$$

Proof: We are allowed to restrict ourselfs to the case $l \ll k$ (more precise: $l < k - 2$), since in the case $l \approx k$ the second derivative of Φ and therefore the Gaussian curvature is comparable on the regions $x \approx 2^k\gamma$ and $x \approx 2^l\gamma$. After rescaling, we are back in the classical case with (almost) constant curvature.
The procedure differs a bit from the previous lemmas: Let $(x, y, z) = \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu$. We claim that the number of tripels (j', l', i') with the property $\xi \in \Gamma_{kj'n} + \Gamma^\delta_{l'i'p}$ is bounded by a fixed constant. Without loss of generality, we may assume $l' \leq l$ and in the case $l' = l$ furthermore $i' \leq i$ by interchanging the parameters if necessary.

Step 1 *If* $(l', i') \neq (l, i)$ *then* $j \leq j'$.

The assumption is not necessary, since in the case $(l', i') = (l, i)$ we may ensure $j \leq j'$ by interchanging the parameters with and without primes.

Case 1: $l' \neq l$

In this case is $l' < l$, i.e. $l' + 1 \leq l$, and therefore

$$\begin{aligned} &(2^k\text{-}1)\gamma + j\gamma_k + (2^l\text{-}1)\gamma \\ \leq &(2^k\text{-}1)\gamma + j\gamma_k + (2^l\text{-}1)\gamma + i\gamma_l \leq x \\ < &(2^k\text{-}1)\gamma + (j'+1)\gamma_k + (2^{l'}\text{-}1)\gamma + (i'+1)\gamma_{l'} \\ \leq &(2^k\text{-}1)\gamma + (j'+1)\gamma_k + (2^{l'+1}\text{-}1)\gamma \\ \leq &(2^k\text{-}1)\gamma + (j'+1)\gamma_k + (2^l\text{-}1)\gamma. \end{aligned}$$

This already implies $j < j' + 1$, i.e. $j \leq j'$.

Case 2: $l' = l$

In this case is $i' < i$, i.e. $i' + 1 \leq i$ and therefore

$$\begin{aligned} &(2^k\text{-}1)\gamma + j\gamma_k + (2^l\text{-}1)\gamma + i\gamma_l \leq x \\ < &(2^k\text{-}1)\gamma + (j'+1)\gamma_k + (2^l\text{-}1)\gamma + (i'+1)\gamma_l \\ \leq &(2^k\text{-}1)\gamma + (j'+1)\gamma_k + (2^l\text{-}1)\gamma + i\gamma_l. \end{aligned}$$

This again implies $j < j' + 1$, i.e. $j \leq j'$.

In the next step, it is usefull to examine the projections in x-$y-$space. Thus we introduce the new curves $\Phi_n(x) = n\beta\Phi\left(\frac{x}{n\beta}\right)$ (analogue Φ_p). Using $\beta < \delta$, it is an easy task to verify

$$\Gamma^\delta_{kjn} = \{(x,y,z) \in \mathbb{R}^3 |\, n\beta \leq z \leq (n+1)\beta,\ \Phi\left(\tfrac{x}{z}\right) \leq \tfrac{y}{z} \leq \Phi\left(\tfrac{x}{z}\right) + \delta,\ x_{kj} \leq x \leq x_{k,j+1}\} \tag{27}$$

$$\subset \tilde{\Gamma}^\delta_{kjn} = \{(x,y,z) \in \mathbb{R}^3 |\, n\beta \leq z \leq (n+1)\beta,\ |y - \Phi_n(x)| \leq 10m\delta,\ x_{kj} \leq x \leq x_{k,j+1}\}, \tag{28}$$

and likewise for Γ^δ_{lip}. Now define the projection $P(x,y,z) = y - x\Phi_n'(x_k)$ on the normal to the graph of Φ_n at point x_k in x-$y-$space.


y
δ
$P(\tilde{\Gamma}^\delta_{lip})$
$\Phi_n = \Phi_p$
$\tilde{\Gamma}^\delta_{kjn}$
$\tilde{\Gamma}^\delta_{lip}$
$\tilde{\Gamma}^\delta_{l'i'p}$
x_k
x
$x_{l'i'}$
x_{li}
x_{kj}
$x_{k,j+1}$

Fig. 3: Projection on the normal vector in the case $n = p$

Additionally, we introduce

$$\xi_{kjn} = (x_{kj}, \Phi_n(x_{kj}), n\beta) \in \Gamma^{\delta}_{kjn} \qquad \xi_{lip} = (x_{li}, \Phi_p(x_{li}), p\beta) \in \Gamma^{\delta}_{lip} \tag{29}$$

and analogue $\xi_{kj'n}$, $\xi_{l'i'p}$.

Step 2

(i) $P(\xi_{l'i'p}) - P(\xi_{lip}) \approx \delta 2^{k(m-1)-l'(\frac{m}{2}-1)}\Delta_{ll'ii'}$

(ii) $P(\xi_{kj'n}) - P(\xi_{kjn}) \gtrsim \delta j(j'-j)$, *falls* $j \leq j'+1$

with $\Delta_{ll'ii'} = 2^{l'(\frac{m}{2}-1)}\dfrac{x_{li}-x_{l'i'}}{\gamma} = 2^{l'(\frac{m}{2}-1)}\left(2^l + i2^{l(1-\frac{m}{2})} - 2^{l'} - i'2^{l'(1-\frac{m}{2})}\right).$

The term $\Delta_{ll'ii'}$ looks somehow artifical. However, we will discover that this quantity is the crucial one, expressing (in some sense) the distance between Γ^{δ}_{lip} and $\Gamma^{\delta}_{l'i'p}$.
Part (i): The mean value theoreme provides the existence of a $\tilde{x} \in (x_{l'i'}, x_{li})$ with

$$\Phi_p(x_{l'i'}) - \Phi_p(x_{li}) = \Phi'_p(\tilde{x})(x_{l'i'} - x_{li}). \tag{30}$$

Thus especially

$$\tilde{x} \leq x_{li} \leq x_{l+1} \underset{l<k-2}{\leq} x_{k-2}$$

and hence

$$\frac{\tilde{x}}{p\beta} \leq \frac{n\beta}{p\beta}\frac{x_{k-2}}{n\beta} \leq \frac{2x_{k-2}}{n\beta} = \frac{2(2^{k-2}-1)\gamma}{n\beta} = \frac{1}{2}\frac{(2^k-4)\gamma}{n\beta} \leq \frac{1}{2}\frac{(2^k-1)\gamma}{n\beta} = \frac{1}{2}\frac{x_k}{n\beta}.$$

Using monotony and convexity of Φ', as well as $\Phi(0) = 0$, we obtain

$$\Phi'\left(\frac{\tilde{x}}{p\beta}\right) \leq \Phi'\left(\frac{1}{2}\frac{x_k}{n\beta}\right) \leq \frac{1}{2}\Phi'\left(\frac{x_k}{n\beta}\right). \tag{31}$$

It follows

$$\begin{aligned}\Phi'_n(x_k) - \Phi'_p(\tilde{x}) =& \Phi'\left(\frac{x_k}{n\beta}\right) - \Phi'\left(\frac{\tilde{x}}{p\beta}\right)\\ \underset{(31)}{\approx}& \Phi'\left(\frac{x_k}{n\beta}\right)\\ \approx& \frac{x_k^{m-1}}{(n\beta)^{m-1}}\\ \approx& \gamma^{m-1}2^{k(m-1)}.\end{aligned} \tag{32}$$

Thus we conclude

$$\begin{aligned}P(\xi_{l'i'p}) - P(\xi_{lip}) =& \Phi_p(x_{l'i'}) - \Phi_p(x_{li}) - (x_{l'i'} - x_{li})\Phi'_n(x_k)\\ \underset{(30)}{=}& (\Phi'_n(x_k) - \Phi'_p(\tilde{x}))(x_{li} - x_{l'i'})\\ =& (\Phi'_n(x_k) - \Phi'_p(\tilde{x}))2^{-l'(\frac{m}{2}-1)}\Delta_{ll'ii'}\gamma\\ \underset{(32)}{\approx}& \gamma^m 2^{k(m-1)}2^{-l'(\frac{m}{2}-1)}\Delta_{ll'ii'}\\ =& \delta 2^{k(m-1)-l'(\frac{m}{2}-1)}\Delta_{ll'ii'}\end{aligned}$$

Part (ii):
Since the projection P depends on k, we are not able to use (i). However, we proceed in a similar manner: Like in part (i), we obtain a $\tilde{x}_k$ between x_{kj} and $x_{kj'}$ with

$$\Phi_n(x_{kj'}) - \Phi_n(x_{kj}) = \Phi_n'(\tilde{x}_k)(x_{kj'} - x_{kj}).$$

Furthermore, we obtain a $\tilde{\tilde{x}}_k$ between $\tilde{x}_k$ and x_k with

$$\begin{aligned} P(\xi_{kj'n}) - P(\xi_{kjn}) =& \Phi_n(x_{kj'}) - \Phi_n(x_{kj}) - (x_{kj'} - x_{kj})\Phi_n'(x_k) \\ =& (\Phi_n'(\tilde{x}_k) - \Phi_n'(x_k))(x_{kj'} - x_{kj}) \\ =& \Phi_n''(\tilde{\tilde{x}}_k)(\tilde{x}_k - x_k)(x_{kj'} - x_{kj}). \end{aligned} \tag{33}$$

Especially $\tilde{\tilde{x}}_k \in (x_k, \tilde{x}_k) \subset (x_k, x_{k+1})$, i.e. $\tilde{\tilde{x}}_k \approx 2^k\gamma$. (34)
In the case $j' \geq j$ we use $\tilde{x}_k > \min\{x_{kj}, x_{kj'}\} = x_{kj} = (2^k\text{-}1)\gamma + j2^{k(1-\frac{m}{2})}\gamma = x_k + j2^{k(1-\frac{m}{2})}\gamma$. (35$a$)

This gives

$$\begin{aligned} P(\xi_{kj'n}) - P(\xi_{kjn}) \; & \underset{(33)}{=} \; \Phi_n''(\tilde{\tilde{x}}_k)(\tilde{x}_k - x_k)(x_{kj'} - x_{kj}) \\ & \underset{(34)}{\approx} \; (2^k\gamma)^{m-2}(\tilde{x}_k - x_k)(x_{kj'} - x_{kj}) \\ & \underset{(35a)}{\geq} \; (2^k\gamma)^{m-2}\; j2^{k(1-\frac{m}{2})}\gamma \; (j'-j)2^{k(1-\frac{m}{2})}\gamma \\ & = \; \gamma^m\, j(j'-j) \\ & = \; \delta j(j'-j). \end{aligned}$$

In the other case $j' < j$ the assumption ensures $j = j'+1$, i.e. $j' - j = -1$. Now we use $\tilde{x}_k < \max\{x_{kj}, x_{kj'}\} = x_{kj} = (2^k\text{-}1)\gamma + j2^{k(1-\frac{m}{2})}\gamma = x_k + j2^{k(1-\frac{m}{2})}\gamma$. (35$b$)
In this case we obtain

$$\begin{aligned} P(\xi_{kj'n}) - P(\xi_{kjn}) \; & \underset{(33)}{=} \; \Phi_n''(\tilde{\tilde{x}}_k)(\tilde{x}_k - x_k)(x_{kj'} - x_{kj}) \\ & \underset{(34)}{\approx} \; (2^k\gamma)^{m-2}(\tilde{x}_k - x_k)(-2^{k(1-\frac{m}{2})}\gamma) \\ & \underset{(35b)}{\geq} \; -(2^k\gamma)^{m-2}\; j2^{k(1-\frac{m}{2})}\gamma \; 2^{k(1-\frac{m}{2})}\gamma \\ & = \; -\gamma^m\, j \\ & = \; -\delta j \;=\; \delta j(j'-j), \end{aligned}$$

completing step 2.
The next task will be to estimate the size of the pieces Γ^δ_{kjn} of our decomposition, with respect to the projection P. The size of a set U with respect to P is measured by $diam_P(U) = \sup\{|P(u) - P(v)| : u, v \in U\}$.

Step 3

(i) $diam_P(\Gamma^\delta_{lip}) \lesssim \delta 2^{k(m-1)-l(\frac{m}{2}-1)}$

(ii) $diam_P(\Gamma^\delta_{kjn}) \lesssim \delta 2^{k\frac{m}{2}}$

Part (i):

$$\begin{aligned} diam_P(\Gamma^\delta_{lip}) =& P(x_{li}, \Phi_p(x_{li}) + \mathcal{O}(\delta), p\beta) - P(x_{l,i+1}, \Phi_p(x_{l,i+1}), p\beta) \\ =& P(x_{li}, \Phi_p(x_{li}), p\beta) - P(x_{l,i+1}, \Phi_p(x_{l,i+1}), p\beta) + \mathcal{O}(\delta) \\ =& P(\xi_{lip}) - P(\xi_{l,i+1,p}) + \mathcal{O}(\delta) \end{aligned}$$

We apply step 2(i), replacing l' by l, i by $i+1$ and i' by i. Then holds $\Delta_{ll,i+1,i} = 2^{l(\frac{m}{2}-1)}\left((i+1)2^{l(1-\frac{m}{2})} - i2^{l(1-\frac{m}{2})}\right) = 1$ and

$$\begin{aligned} diam_P(\Gamma^\delta_{lip}) =& P(\xi_{lip}) - P(\xi_{l,i+1,p}) + \mathcal{O}(\delta) \\ \approx& \delta 2^{k(m-1)-l(\frac{m}{2}-1)}\Delta_{ll,i+1,i} + \mathcal{O}(\delta) \\ =& \delta 2^{k(m-1)-l(\frac{m}{2}-1)} + \mathcal{O}(\delta) \\ \overset{l\leq k}{\lesssim}& \delta 2^{k(m-1)-l(\frac{m}{2}-1)}. \end{aligned}$$

Part (ii):

$$\begin{aligned} diam_P(\Gamma^\delta_{kjn}) =& P(x_{k,j+1}, \Phi_n(x_{k,j+1}) + \mathcal{O}(\delta), n\beta) - P(x_{kj}, \Phi_n(x_{kj}), n\beta) \\ =& P(x_{k,j+1}, \Phi_n(x_{k,j+1}), n\beta) - P(x_{kj}, \Phi_n(x_{kj}), n\beta) + \mathcal{O}(\delta) \\ =& P(\xi_{k,j+1,n}) - P(\xi_{kjn}) + \mathcal{O}(\delta) \end{aligned}$$

Step 2(ii) implies

$$P(\xi_{kj'n}) - P(\xi_{kjn}) \gtrsim \delta j(j'-j), \qquad \text{if } j \leq j'+1$$

or, equivalently

$$P(\xi_{kjn}) - P(\xi_{kj'n}) \lesssim \delta j(j-j'), \qquad \text{if } j \leq j'+1,$$

so especially

$$P(\xi_{k,j+1n}) - P(\xi_{kjn}) \lesssim (j+1)\delta \leq \delta 2^{k\frac{m}{2}}.$$

This leads to

$$\begin{aligned} diam_P(\Gamma^\delta_{kjn}) =& P(\xi_{k,j+1,n}) - P(\xi_{kjn}) + \mathcal{O}(\delta) \\ \lesssim& \delta 2^{k\frac{m}{2}} + \mathcal{O}(\delta) \\ \lesssim& \delta 2^{k\frac{m}{2}}. \end{aligned}$$

A further step will analyse the quantity $\Delta_{ll'ii'}$. Here we exploit $\Gamma^\delta_{kj'n} + \Gamma^\delta_{l'i'p} \neq \emptyset$.

Step 4 $\Delta_{ll'ii'} \lesssim 1$

We apply lemma 7.5. Since

$$\xi_\alpha \in \Gamma^\delta_\alpha, \qquad \alpha = (k,j,n),(k,j',n),(l,i,p),(l',i',p)$$

the lemma yields

$$\begin{aligned} & P(\xi_{kj'n}) - P(\xi_{kjn}) + P(\xi_{l'i'p}) - P(\xi_{lip}) \\ \leq \; & diam_P(\Gamma^\delta_{kjn}) + diam_P(\Gamma^\delta_{kj'n}) + diam_P(\Gamma^\delta_{lip}) + diam_P(\Gamma^\delta_{l'i'p}). \end{aligned} \tag{36}$$

In step 1, we established $j \leq j'$ (at least if $(l,i) \neq (l',i')$, but in the case $(l,i) = (l',i')$ we may also assume this). The application of step 2 and step 3 results in

$$\begin{aligned} \delta 2^{k(m-1)-l'(\frac{m}{2}-1)}\Delta_{ll'ii'} \leq & \delta 2^{k(m-1)-l'(\frac{m}{2}-1)}\Delta_{ll'ii'} + \delta j(j'-j) \\ & \overset{(36)}{\lesssim} \delta 2^{k\frac{m}{2}} + \delta 2^{k(m-1)-l(\frac{m}{2}-1)} + \delta 2^{k(m-1)-l'(\frac{m}{2}-1)} \end{aligned} \tag{37}$$

Now we use $l' \leq l$ and $l' < k$, since the last one implies $k(m-1) - l'(\frac{m}{2}-1) = k\frac{m}{2} + (k-l')(\frac{m}{2}-1) \geq k\frac{m}{2}$. Thus (37) transformes into

$$2^{k(m-1)-l'(\frac{m}{2}-1)}\Delta_{ll'ii'} \lesssim 2^{k(m-1)-l(\frac{m}{2}-1)} + 2^{k(m-1)-l'(\frac{m}{2}-1)} + 2^{k\frac{m}{2}} \lesssim 2^{k(m-1)-l'(\frac{m}{2}-1)},$$

i.e. $\Delta_{ll'ii'} \lesssim 1$.
The former results will be merged into the following step, which states that there are not „too many“ (l',i') appropriate to a given (l,i).

Step 5 *One of these three alternatives holds:*

(i) $l' = l$*: Then* $|i-i'| \lesssim 1$.

(ii) $l' = l-1$*: Then* $i \lesssim 1$ *and* $2^{l'\frac{m}{2}} - i' \lesssim 1$.

(iii) $l' < l-1$*: Then* $l', l \lesssim 1$ *and especially* $i', i \lesssim 1$.

Especially $l' \approx l$ *is necessary.*

Let me remind you that we assumed $l' \leq l$ and even $i' \leq i$ for $l' = l$. The proof is divided into three cases:
Case 1: $l' = l$
According to step 4, we know

$$1 \gtrsim \Delta_{llii'} = 2^{l(\frac{m}{2}-1)}\left(2^l + i2^{l(1-\frac{m}{2})} - 2^l - i'2^{l(1-\frac{m}{2})}\right) = i - i' \geq 0.$$

Case 2: $l' = l-1$
Step 4 now reads

$$1 \gtrsim \Delta_{ll'ii'} = 2^{l'(\frac{m}{2}-1)}\left(2^{l'+1} + i2^{(l'+1)(1-\frac{m}{2})} - 2^{l'} - i'2^{l'(1-\frac{m}{2})}\right) = \underbrace{2^{l'\frac{m}{2}} - i'}_{\geq 0} + \underbrace{i2^{1-\frac{m}{2}}}_{\geq 0},$$

hence

$$1 \gtrsim i2^{1-\frac{m}{2}} \approx i \qquad \text{and} \qquad 1 \gtrsim 2^{l'\frac{m}{2}} - i'.$$

Case 3: $l > l' + 1$
In this case, step 4 yields

$$\begin{aligned} 1 \gtrsim \Delta_{ll'ii'} =& 2^{l'(\frac{m}{2}-1)} \left(2^l + i2^{l(1-\frac{m}{2})} - 2^{l'} - i'2^{l'(1-\frac{m}{2})}\right) \\ \gtrsim& 2^{l'(\frac{m}{2}-1)} \left(2^l - 2^{l'+1}\right) \gtrsim 2^{l'(\frac{m}{2}-1)}2^l \geq 2^l, \end{aligned}$$

which already implies $l' \leq l \lesssim 1$ and thus $i \leq 2^{l\frac{m}{2}} \lesssim 1$, as well as $i' \lesssim 1$.


y
δ
$Q(\tilde{\Gamma}^\delta_{kjn})$
$\tilde{\Gamma}^\delta_{kjn}$
$\Phi_n = \Phi_p$
$\tilde{\Gamma}^\delta_{l'i'p}$
$\tilde{\Gamma}^\delta_{lip}$
x
$x_{l'i'}$
x_{li}
x_{kj}
$x_{k,j+1}$

Fig. 4: Projection on the y-axis in the case $n = p$

To come to a similar conclusion concerning j and j', we consider the projection on the $y-$axis, denoted $Q(x, y, z) = y$. Then -cf. (29)- $Q(\xi_{kjn}) = \Phi_n(x_{kj})$ and $Q(\xi_{lip}) = \Phi_p(x_{li})$, analogue for the primed coordinates.

Step 6

(i) $Q(\xi_{lip}) - Q(\xi_{l'i'p}) \lesssim \delta 2^{l\frac{m}{2}}$

(ii) $Q(\xi_{kj'n}) - Q(\xi_{kjn}) \approx \delta 2^{k\frac{m}{2}}(j' - j)$

Part (ii): Again, we use the mean value theorem: There exists a $\tilde{x}$ between x_{kj} and $x_{kj'}$ fulfilling

$$Q(\xi_{kj'n}) - Q(\xi_{kjn}) = \Phi_n(x_{kj'}) - \Phi_n(x_{kj}) = \Phi_n'(\tilde{x})(x_{kj'} - x_{kj}) = \Phi'\left(\frac{\tilde{x}}{n\beta}\right)(x_{kj'} - x_{kj}).$$

Notice that $\tilde{x} \approx x_k \approx 2^k\gamma$, i.e.

$$\begin{aligned} Q(\xi_{kj'n}) - Q(\xi_{kjn}) \approx& (2^k\gamma)^{m-1}(x_{kj'} - x_{kj}) \\ =& 2^{k(m-1)}\gamma^{m-1}(j' - j)2^{k(1-\frac{m}{2})}\gamma \\ =& \delta 2^{k\frac{m}{2}}(j' - j). \end{aligned}$$

Part (i): In this case, mean value theorem provides a $\bar{x} \approx x_l \approx 2^l\gamma$ such that

$$\begin{aligned} Q(\xi_{lip}) - Q(\xi_{l'i'p}) =& \Phi_p(x_{li}) - \Phi_p(x_{l'i'}) \\ =& \Phi_p'(\bar{x})(x_{li'} - x_{l'i'}) \\ \approx& (2^l\gamma)^{m-1}\ 2^{-l'(\frac{m}{2}-1)}\gamma\Delta_{ll'ii'} \\ =& \delta\ 2^{l\frac{m}{2}}\ 2^{(l-l')(\frac{m}{2}-1)}\ \Delta_{ll'ii'}. \end{aligned}$$

Since $\Delta_{ll'ii'} \lesssim 1$ by step 4 and $l' \approx l$ according to 5, the equation transfares into

$$Q(\xi_{lip}) - Q(\xi_{l'i'p}) \lesssim \delta 2^{l\frac{m}{2}},$$

thus completing step 6.

Step 7 $|j - j'| \lesssim 1$

We have

$$\begin{aligned}
diam_Q(\Gamma^\delta_{kjn}) &= \sup_{\xi,\eta\in\Gamma^\delta_{kjn}} |Q(\xi) - Q(\eta)| \\
&= Q(x_{kj}, \Phi_n(x_{kj}) + \mathcal{O}(\delta), n\beta) - Q(x_{k,j-1}, \Phi_n(x_{k,j-1}), n\beta) \\
&= \Phi_n(x_{kj}) + \mathcal{O}(\delta) - \Phi_n(x_{k,j-1}) \\
&\underset{Step\ 6(ii)}{\approx} \delta 2^{k\frac{m}{2}} + \mathcal{O}(\delta) \\
&\lesssim \delta 2^{k\frac{m}{2}}.
\end{aligned}$$

Using step 6(i) we observe $diam_Q(\Gamma^\delta_{lip}) \lesssim \delta 2^{l\frac{m}{2}} \leq \delta 2^{k\frac{m}{2}}$ and $diam_Q(\Gamma^\delta_{l'i'p}) \lesssim \delta 2^{k\frac{m}{2}}$. Now we again apply lemma 7.5:

$$\begin{aligned}
Q(\xi_{kj'n}) - Q(\xi_{kjn}) &+ Q(\xi_{l'i'p}) - Q(\xi_{lip}) \\
&\leq diam_Q(\Gamma^\delta_{kj'n}) + diam_Q(\Gamma^\delta_{kjn}) + diam_Q(\Gamma^\delta_{l'i'p}) + diam_Q(\Gamma^\delta_{lip}) \\
&\lesssim \delta 2^{k\frac{m}{2}}.
\end{aligned}$$

It follows

$$\begin{aligned}
\delta 2^{k\frac{m}{2}}(j' - j) &\underset{Schritt\ 6(ii)}{\approx} Q(\xi_{kj'n}) - Q(\xi_{kjn}) \\
&\lesssim Q(\xi_{lip}) - Q(\xi_{l'i'p}) + \delta 2^{k\frac{m}{2}} \\
&\underset{Schritt\ 6(i)}{\lesssim} \delta 2^{l\frac{m}{2}} + \delta 2^{k\frac{m}{2}} \\
&\lesssim \delta 2^{k\frac{m}{2}},
\end{aligned}$$

and thus $j' - j \lesssim 1$. This already implies the desired estimate, since $j \leq j' + 1$ according to step 1.
Togehter with step 5, step 7 completes the proof of lemma 4.3.

4.3 Summary of the results

Corollary 4.4
For all $\xi \in \mathbb{R}^3$, we have

$$\#\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} \leq 18C\beta^{-1}, \tag{38}$$

where C is the constant from lemma 4.3.

Proof: Let

$$\begin{aligned} M =&\{(n,p)|\exists k,l,j,i : \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\},\\ M(n,p) =&\{k|\exists l,j,i : \xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \leq k\},\\ M(n,p;k) =&\{(l,j,i)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\}. \end{aligned}$$

Then the previous lemma 4.1, 4.2 and 4.3 states

$$\#M \leq 3\beta^{-1} \tag{39}$$

$$\#M(n,p) \leq 3 \qquad \forall n,p \tag{40}$$

$$\#M(n,p;k) \leq C \qquad \forall n,p,k. \tag{41}$$

We procced by

$$\begin{aligned} \{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} =&\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \leq k\} \cup \{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \geq k\}\\ =&\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \leq k\} \cup \{(\alpha,\mu)|\xi \in \Gamma^\delta_\mu + \Gamma^\delta_\alpha,\ k \leq l\}, \end{aligned}$$

and

$$\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \leq k\} = \{(\alpha,\mu)|(n,p) \in M,\ k \in M(n,p),\ (l,j,i) \in M(n,p;k)\}.$$

It follows

$$\begin{aligned} \#\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu\} \leq& 2\#\{(\alpha,\mu)|\xi \in \Gamma^\delta_\alpha + \Gamma^\delta_\mu,\ l \leq k\}\\ =&2\sum_{(n,p)\in M}\ \sum_{k\in M(n,p)} \#M(n,p;k)\\ \leq& 2 \cdot 3\beta^{-1} \cdot 3 \cdot C = 18C\beta^{-1} \end{aligned}$$

using Fubini's theorem. □

If we enlarge the sets Γ^δ_α in the right manner, the statemant essentialy remains valid:

Corollary 4.5
For the sets Γ^δ_{kjn}, we introduce their adjacent sets

$$\Gamma^\delta_{kjn}(u,v,w) = (0, v\delta, 0) + \Gamma^\delta_{k,j+u,n+w}, \qquad u,v,w \in \{-1,0,1\},$$

as well as their „doubling“

$$G^\delta_{kjn} = \bigcup_{u,v,w\in\{-1,0,1\}} \Gamma^\delta_{kjn}(u,v,w).$$

Then for every $\xi \in \mathbb{R}^3$:

$$\#\{(\alpha,\mu)|\xi \in G^\delta_\alpha + G^\delta_\mu\} \leq 3^3 \cdot 18C\beta^{-1}.$$

The last variation of these lemma is the version we want to apply finally.

Corollary 4.6

Let $1 \leq s < \infty$ *and* G^{δ}_{kjn} *like above. Then there exists a constant* $C_s > 0$ *satisfying the following: Let* $f_{\alpha\mu} \in C^{\infty}_{0}(\mathbb{R}^3)$*,* $k, l = 0, \ldots, |\log_2 \gamma| - 1$*,* $j = 0, \ldots 2^{k\frac{m}{2}} - 1$*,* $i = 0, \ldots 2^{l\frac{m}{2}} - 1$*,* $n, p = \frac{1}{\beta}, \ldots, \frac{2}{\beta} - 1$*, be non-negative functions fulfilling*

$$\mathit{supp}\, f_{\alpha\mu} \subset G^{\delta}_{\alpha} + G^{\delta}_{\mu}$$

Then for all $\xi \in \mathbb{R}^3$

$$\left(\sum_{\alpha\mu} f_{\alpha\mu}(\xi) \right)^{s} \leq C_s \beta^{1-s} \sum_{\alpha\mu} |f_{\alpha\mu}(\xi)|^{s}.$$

5 L_p-estimates for convolutions

Consider two cuboids in $\mathbb{R}^3$ with two short and one long edge. Both cuboids shall lay at parallel planes (we will concretize this soon). If we form the convolution of two functions, each one supported in one of the cuboids, what can we say about the L_p-Norm, depending on size and relative position?

5.1 Introduction to the Problem and formulation of the result

For $i = 1, 2$ let $A_i = \{(x, y) : |x| \le \frac{\gamma_i}{2},\ |y - m_i x| \le \frac{\delta}{2}\}$. We assume that $\delta \ll \gamma_2, \gamma_1$ and moreover, that the slope m_i of the boxes is bounded by an absolute constant (i.e. not depending on δ and γ_i). For convenience, let $\gamma_2 \le \gamma_1$.



Fig.5: parallelogram A_i

Furthermore, we configure the position of the figures relative to each other. Let $\alpha = \angle(A_1, A_2)$ be the angle between the parallelograms, i.e. between their longer sides. The assumption of the boundedness of the slopes guarantees $\sin\alpha \approx \alpha \approx \tan\alpha$. (42)



Fig.6: parallelograms A_i

Now we introduce the paralleloids $Q_i = A_i \times (0,\beta) \subset \mathbb{R}^3$. Let $\xi_i \in \mathbb{R}^3$ and ϕ_i a bump function adopted to $\xi_i + Q_i$.
The main result of this chapter is the following:

Lemma 5.1
Let $1 \leq s < \infty$. Then

$$\int_{\mathbb{R}^3} |\phi_1 * \phi_2|^s \, \mathrm{d}x \lesssim \frac{(\beta\delta)^{(s+1)}}{(1+\frac{\gamma_2}{\delta}\alpha)^{s-1}}\gamma_2^s\gamma_1. \tag{43}$$

5.2 Proof

At the beginning a simple remark:

Remark 5.2
For $C > 0$ and $i = 1,2$ let B_i be symmetric ($B_i = -B_i$) subsets of $\mathbb{R}^n$, $x_i \in \mathbb{R}^n$, supp $\psi_i \subset x_i + B_i$ and $\|\psi_i\|_\infty \leq C$. Then

$$\|\psi_1 * \psi_2\|_\infty \leq C^2 |B_1 \cap \sup_z (z + B_2)|. \tag{44}$$

Proof:

$$\begin{aligned}|\psi_1 * \psi_2|(x) =& \Big|\int\limits_{\mathbb{R}^n} \psi_1(y)\psi_2(x-y) \, \mathrm{d}y\Big| \\ \leq& \int\limits_{\mathbb{R}^n} |\psi_1(x_1+y)\psi_2(x-x_1-y)| \, \mathrm{d}y\end{aligned}$$

If the integrand does not vanish, we have $x_1 + y \in supp\,\psi_1 \subset x_1 + B_1$, i.e. $y \in B_1$. Furthermore must $x - x_1 - y \in supp\,\psi_2 \subset x_2 + B_2$ be valid, i.e. $y \in x - x_1 - x_2 - B_2 = x - x_1 - x_2 + B_2$. We conclude

$$\begin{aligned}|\psi_1 * \psi_2|(x) \leq & \int\limits_{B_1\cap(x-x_1-x_2+B_2)} |\psi_1(x_1+y)\psi_2(x-x_1-y)|\,\mathrm{d}y \\ \leq & C^2|B_1 \cap (x - x_1 - x_2 + B_2)| \\ \leq & C^2 \sup_z |B_1 \cap (z + B_2)|\end{aligned}$$

as claimed. □

Remark

If the sets lay like in figure 6, the supremum is achieved (amongst others) in $z = 0$, as we can see clearly in the picture. In this concrete situation, a formal proof can be obtained with some efforts by an elementary computation.

However, it would be more typical for a mathematician to conjecture a more general statement: „Let K and L be two convex bodies in $\mathbb{R}^n$ symmmetric with respect to the origin (i.e. $K = -K$). Then for all $z \in \mathbb{R}^n$ follows $|K \cap (z + L)| \leq |K \cap L|$.“

As it is often in convex geometry, the statement is quite easy to understand, whereas a proof is not immediatly clear. One might use the inequality of Brunn-Minkowski (cf. lemma 7.4).

Anyway, we now need to estimate $|A_1 \cap A_2|$:

Lemma 5.3

We have

$$|A_1 \cap A_2| \lesssim \frac{\delta^2\gamma_2}{\max\{\delta, \gamma_2\alpha\}} \approx \frac{\delta^2\gamma_2}{\delta + \gamma_2\alpha}.$$

Proof: Enlongating the two parallelograms in direction of their longer edges to infinity, we obtain by a elementary computation (cf. figure 7)

$$|A_1 \cap A_2| \leq \frac{\delta^2}{\sin\alpha} \overset{(42)}{\approx} \frac{\delta^2}{\alpha}. \tag{45}$$



Fig. 7: $A_1 \cap A_2$

This expression is indeed to large for small angles; in this case we therefore apply the trivial estimate

$$|A_1 \cap A_2| \leq |A_2| = \delta\gamma_2. \tag{46}$$

The changeover appears if the corners of the small parallelogramm get into touch with the large parallelogramm, i.e. if $\alpha \approx \frac{\delta}{\gamma_2}$. Given $\gamma_2\alpha \geq \delta$ we thus apply (45) and find

$$|A_1 \cap A_2| \lesssim \frac{\delta^2}{\alpha} = \frac{\delta^2\gamma_2}{\max\{\delta, \gamma_2\alpha\}}.$$

In the contrary case $\gamma_2\alpha < \delta$ we apply (46) and find

$$|A_1 \cap A_2| \leq \delta\gamma_2 = \frac{\delta^2\gamma_2}{\max\{\delta, \gamma_2\alpha\}},$$

and thus the desired estimate in both cases. □

This lemma can be proved in a similar manner, as basic as before:

Lemma 5.4

$$|A_1 + A_2| \lesssim \gamma_1(\delta + \gamma_2\alpha).$$

Now we are able to start with the proof of lemma 5.1:

Proof:

$$\begin{aligned}
\int_{\mathbb{R}^3} |\phi_1 * \phi_2|^s \,\mathrm{d}x \quad & \leq \quad \|\phi_1 * \phi_2\|_\infty^s \; |\,supp(\phi_1 * \phi_2)| \\
& \underset{\text{Lemma 5.2}}{\lesssim} \quad \Big(\sup_{\eta \in \mathbb{R}^2 \times \mathbb{R}} |Q_1 \cap (\eta + Q_2)| \Big)^s \; |\xi_1 + Q_1 + \xi_2 + Q_2| \\
& = \quad |Q_1 \cap Q_2|^s \; |Q_1 + Q_2| \\
& = \quad |\big(A_1 \cap A_2\big) \times (0,\beta)|^s \; |\big(A_1 \times (0,\beta)\big) + \big(A_2 \times (0,\beta)\big)| \\
& = \quad |A_1 \cap A_2|^s \beta^s \; |\big(A_1 + A_2\big) \times (0, 2\beta)| \\
& \underset{\text{Lemma 5.3}}{\lesssim} \quad \left(\frac{\delta^2\gamma_2}{\delta + \gamma_2\alpha} \right)^s \; |\big(A_1 + A_2\big)| \; \beta^{s+1} \\
& \underset{\text{Lemma 5.4}}{\lesssim} \quad \left(\frac{\delta^2\gamma_2}{\delta + \gamma_2\alpha} \right)^s \; \gamma_1(\delta + \gamma_2\alpha) \; \beta^{s+1} \\
& = \quad \frac{\delta^{2s}\, \gamma_2^s \, \gamma_1}{(\delta + \gamma_2\alpha)^{s-1}} \; \beta^{s+1} \\
& = \quad \frac{(\beta\delta)^{s+1}\, \gamma_2^s \, \gamma_1}{(1 + \frac{\gamma_2}{\delta}\alpha)^{s-1}}.
\end{aligned}$$

6 Proof of theorem 2.2

Befor we start to complete the main proof, we need a further lemma, which concretises the general results from the previous chapter in our special situation.

6.1 Application of the results from chapter 5

LEMMA 6.1
Let $1 \leq s < \infty$. The functions ϕ_{kjn} introduced in the previous chapter satisfy the following estimates:

(i) If $|k-l| \gg 1$ then

$$\int |\phi_{kjn} * \phi_{lip}|^s \, \mathrm{d}x \lesssim (\beta\delta\gamma)^{s+1} 2^{\frac{k+l}{2}(1+s-sm)} \; 2^{-\frac{|k-l|}{2}(m-1)(s-1)}. \tag{47}$$

(ii) If $|k-l| \lesssim 1$, we find either a function $f : I_k \to I_l$ with $\forall i \in I_l : |\{j : f(j) = i\}| \lesssim 1$ or a function $g : I_l \to I_k$ with $\forall j \in I_k : |\{i : g(i) = j\}| \lesssim 1$ such that

$$\int |\phi_{kjn} * \phi_{lip}|^s \, \mathrm{d}x \lesssim (\beta\delta\gamma)^{s+1} 2^{\frac{k+l}{2}(1+s-sm)} \left[\frac{2^{\frac{k+l}{2}\frac{m}{2}}}{1+h(i,j)} \right]^{s-1}, \tag{48}$$

where

$$h(i,j) = |f(j) - i| \qquad \text{or} \qquad h(i,j) = |j - g(i)|. \tag{49}$$


y
$P_{xy}(\tilde{\Gamma}^{\delta}_{kjn})$
x
$P_{xy}(\tilde{\Gamma}^{\delta}_{lip})$

Fig.8: Angle between the translations to the origin of the projections

Proof: Recall the definition of $\tilde{\Gamma}^{\delta}_{kjn} = \{(x,y,z) \in \mathbb{R}^3 | \, n\beta \leq z \leq (n+1)\beta, \; |y - \Phi_n(x)| \leq 10m\delta, \; x_{kj} \leq x \leq x_{k,j+1}\}$. The projections of these sets on x-$y-$space are contained in parallelograms with slope $\Phi'_n(x_{kj})$, thickness $\mathcal{O}(\delta)$ and width γ_{kj}. When we shift their centers to the origin, they intersect with angle

$$\alpha = \angle(\tilde{\Gamma}^{\delta}_{kjn}, \tilde{\Gamma}^{\delta}_{lip}) = |\arctan \Phi'_n(x_{kj}) - \arctan \Phi'_p(x_{li})|. \tag{50}$$

Since Φ' and therefore as well $\Phi'_n = \Phi'\left(\frac{\cdot}{n\beta}\right)$ and Φ'_p are bounded from above and from below independly of n and p, we have

$$\alpha \approx |\Phi'_n(x_{kj}) - \Phi'_p(x_{li})|. \tag{51}$$

At first we consider the case (i) $|k-l| \gg 1$ and assume without loss of generality $l \leq k$, we obtain due to the dyadic nature of the construction

$$\begin{aligned}\alpha \approx & \Phi'_n(x_{kj}) \approx (2^k\gamma)^{m-1} \\ = & \frac{\gamma^m}{2^{k(1-\frac{m}{2})}\gamma} 2^{k\frac{m}{2}} \\ = & \frac{\delta}{\gamma_k} 2^{k\frac{m}{2}}.\end{aligned} \tag{52}$$

The case (ii) $|k-l| \lesssim 1$ is a bit more complicated. This is due to the fact that we decomposed orthogonal to the $x-$axis, regardless the cone-like shape of the surface. Formula (51) illustrates the difficulies: In the special case $n = p$ we get rid of the difference by the fundamental theorem of calculus, but unfortunatly, the general case appears much harder.

Instead of this, we choose an other approach: Let k, l, n, p be fixed and $a_i = \Phi'_p(x_{li})$, $b_j = \Phi'_n(x_{kj})$. Then for all j, j'

$$\begin{aligned}|b_j - b_{j'}| &= |\Phi'_n(x_{kj}) - \Phi'_n(x_{kj'})| \\ &\approx \Phi''(x_k)|x_{kj} - x_{kj'}| \\ &\approx (2^k\gamma)^{m-2}\gamma_k|j-j'| \\ &= 2^{k(\frac{m}{2}-1)}\gamma^{m-1}|j-j'| \\ &= \frac{\delta}{\gamma_k}|j-j'|.\end{aligned} \tag{53}$$

In the same manner we obtain

$$|a_i - a_{i'}| \approx \frac{\delta}{\gamma_l}|i-i'| \overset{|k-l|\lesssim 1}{\approx} \frac{\delta}{\gamma_k}|i-i'|. \tag{54}$$

This basicly means that we have a good idea how to compare the a_i with each other, the same with the $b'_j s$. But we lack control in comparing some a_i with a b_j. Therefore we apply the abstract result of lemma 7.6. At first we check the preconditions:

The sequences $a_i, i \in I_l$ and $b_j, j \in I_k$ are increasing, since (for instance) $x_{li} < x_{li} + \gamma_l = x_{l,i+1}$ and Φ'_p is monotonously increasing. Thus we may apply lemma 7.6 to the renormed sequences $\frac{\gamma_k}{\delta}a_i$ and $\frac{\gamma_k}{\delta}b_j$. Provided $b_{2^{km/2}-1} - b_0 \leq a_{2^{lm/2}-1} - a_0$ we get a function $f : I_k \to I_l$ almost injectiv, i.e. $\forall i \in I_l : |\{j \in I_k : f(j) = i\}| \lesssim 1$, and fulfilling

$$|a_i - b_j| \geq \frac{1}{2}|a_i - a_{f(j)}| \quad \forall i \in I_l \forall j \in I_k. \tag{55}$$

Consequently

$$\begin{aligned}\alpha \approx &|\Phi_n'(x_{kj}) - \Phi_p'(x_{li})|\\ =&|b_j - a_i|\\ \overset{(55)}{\gtrsim}& |a_i - a_{f(j)}|\\ \overset{(54)}{\approx}& \frac{\delta}{\gamma_k}|i - f(j)|.\end{aligned}$$

Provided $b_{2^{km/2}-1} - b_0 > a_{2^{lm/2}-1} - a_0$ we switch the roles of a_i and b_j and obtain a correspondingly result; in any case, we obtain a function h as desired and fulfilling

$$\alpha \gtrsim \frac{\delta}{\gamma_k} h(i,j). \tag{56}$$

Consolidation of (52) and (56) yields

$$\frac{\gamma_k}{\delta}\alpha = \begin{cases} 2^{k\frac{m}{2}}, & \text{falls } l \ll k \\ h(i,j), & \text{falls } |k-l| \lesssim 1. \end{cases} \tag{57}$$

Using lemma 5.1 and taking into account $l \leq k$, i.e. $\gamma_k \leq \gamma_l$, we conclude

$$\begin{aligned}\int |\phi_{kjn} * \phi_{lip}|^s \,\mathrm{d}x \lesssim &\frac{(\beta\delta)^{s+1}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}}\gamma_k^s\gamma_l\\ =&\frac{(\beta\delta\gamma)^{s+1}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}}2^{k(1-\frac{m}{2})s}2^{l(1-\frac{m}{2})}\\ =&\frac{(\beta\delta\gamma)^{s+1}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}}2^{\frac{k}{2}(2s-ms)}2^{\frac{l}{2}(2-m)}\\ =&(\beta\delta\gamma)^{s+1}2^{\frac{k+l}{2}(1+s-sm)} \cdot \frac{2^{\frac{k}{2}(s-1)+\frac{l}{2}(ms-m-s+1)}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}}.\end{aligned}$$

It remains to consider the second expression. In the case $l \ll k$, it is transformed by (57) into

$$\begin{aligned}\frac{2^{\frac{k}{2}(s-1)+\frac{l}{2}(ms-m-s+1)}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}} \approx &\frac{2^{\frac{k}{2}(s-1)+\frac{l}{2}(ms-m-s+1)}}{2^{k\frac{m}{2}(s-1)}}\\ =&2^{\frac{k}{2}(s-1-ms+m)+\frac{l}{2}(ms-m-s+1)}\\ =&2^{-\frac{k-l}{2}(m-1)(s-1)},\end{aligned}$$

whereas in the case $|k-l| \lesssim 1$, it gives

$$\begin{aligned}\frac{2^{\frac{k}{2}(s-1)+\frac{l}{2}(ms-m-s+1)}}{(1+\frac{\gamma_k}{\delta}\alpha)^{s-1}} =&\frac{2^{\frac{k}{2}(s-1)+\frac{l}{2}(ms-m-s+1)}}{(1+h(i,j))^{s-1}}\\ \overset{k\approx l}{\approx}&\frac{2^{\frac{k+l}{4}(s-1)+\frac{k+l}{4}(ms-m-s+1)}}{(1+h(i,j))^{s-1}}\\ =&\frac{2^{\frac{k+l}{4}(ms-m)}}{(1+h(i,j))^{s-1}}\\ =&\left(\frac{2^{\frac{k+l}{2}\frac{m}{2}}}{1+h(i,j)}\right)^{s-1},\end{aligned}$$

whereby the claim is verified. □

6.2 Completing the proof

One further intermediate step will be helpfull since we get rid of the split-up in the two different cases from the previous lemma.

Lemma 6.2
Let $\frac{3}{p'} < \frac{1}{q} < \frac{1}{2} + \frac{1}{p'}$ *and* $\frac{1}{s} + \frac{2}{p'} = 1.$ (58)
Then for all k, l, n, p *and for all finite sequences* $a \in \mathbb{R}^{I_k}, b \in \mathbb{R}^{I_l}$ *we have*

$$\begin{aligned}&\sum_{ij} |a_j|^s |b_i|^s \int |\phi_{kjn} * \phi_{lip}|^s \,\mathrm{d}x\\ \lesssim&(\beta\delta\gamma)^{s+1} \|a\|_{q'}^s \|b\|_{q'}^s 2^{\frac{k+l}{2}(1+s-sm+m\left(1-\frac{s}{q'}\right))} 2^{-\frac{|k-l|}{2}(m-1)(s-1)}.\end{aligned}$$

Proof: At first we consider the case $|k-l| \gg 1$. Hölders inequality implies

$$\begin{aligned}\sum_j |a_j|^s \leq \left(\sum_j |a_j|^{q'}\right)^{\frac{s}{q'}} \left(\sum_{j=0}^{2^{k\frac{m}{2}}-1}\right)^{1-\frac{s}{q'}}\\ =\|a\|_{q'}^s 2^{\frac{k}{2}m\left(1-\frac{s}{q'}\right)},\end{aligned}$$

which, together with lemma 6.1, results in the desired estimate.

In the case $|k-l| \leq 1$ we apply lemma 7.1 with parameter $r = \frac{q'}{s}$. (59)
Since we assumed $\frac{1}{q} < \frac{1}{2} + \frac{1}{p'}$, it follows

$$\frac{1}{q'} = 1 - \frac{1}{q} > \frac{1}{2} - \frac{1}{p'} = \frac{1}{2}\left(1 - \frac{2}{p'}\right) \overset{(58)}{=} \frac{1}{2s}, \qquad \text{thus } r = \frac{q'}{s} < 2. \tag{60}$$

Furthermore, we have

$$\begin{aligned} \frac{1}{r'} =& 1-\frac{s}{q'} = 1-s+\frac{s}{q} \\ >& 1-s+s\frac{3}{p'} = 1-s+\frac{3s}{2}\frac{2}{p'} \\ \overset{(58)}{=}& 1-s+\frac{3s}{2}\left(1-\frac{1}{s}\right) = -(s-1)+\frac{3}{2}(s-1) \\ =& \frac{s-1}{2}, \end{aligned}$$

i.e.

$$Q := (s-1)\frac{r'}{2} < 1. \tag{61}$$

Assume for simplicity once more that $l \leq k$. To apply the lemma, we have to analyse the kernel $G_{\frac{k+l}{2}}(x) = \frac{2^{\frac{k+l}{2}\frac{m}{2}}}{1+x}$ aus (48), $x \in [0, 2^{k\frac{m}{2}}-1]$. It fulfills

$$\begin{aligned} \int_0^{2^{k\frac{m}{2}}-1} G_{\frac{k+l}{2}}^Q \, \mathrm{d}x \leq \int_1^{2^{k\frac{m}{2}}} \left(\frac{2^{k\frac{m}{2}}}{x}\right)^Q \, \mathrm{d}x = 2^{k\frac{m}{2}} \int_{2^{-k\frac{m}{2}}}^1 x^{-Q} \, \mathrm{d}x \\ \leq 2^{k\frac{m}{2}} \left[\frac{x^{1-Q}}{1-Q}\right]_0^1 \approx 2^{k\frac{m}{2}} \overset{|k-l|\lesssim 1}{\approx} 2^{\frac{k+l}{2}\frac{m}{2}}, \end{aligned}$$

i.e.

$$\|G_{\frac{k+l}{2}}^{s-1}\|_{\frac{r'}{2}} = \|G_{\frac{k+l}{2}}\|_Q^{s-1} \lesssim 2^{\frac{k+l}{2}\frac{m}{2}\frac{s-1}{Q}} \overset{(61)}{\leq} 2^{\frac{k+l}{2}\frac{m}{r'}} \overset{(59)}{=} 2^{\frac{k+l}{2}m\left(1-\frac{s}{q'}\right)}. \tag{62}$$

Next we again apply lemma 6.1, but this time part (ii), though we just discuss the first case $h(i,j) = |f(j)-i|$. The lemma states

$$\begin{aligned} &\sum_{ij} |a_j|^s |b_i|^s \int |\phi_{kjn} * \phi_{lip}|^s \, \mathrm{d}x \\ \lesssim& (\beta\delta\gamma)^{s+1} \sum_{ij} |a_j|^s |b_i|^s 2^{\frac{k+l}{2}(1+s-sm)} \left[\frac{2^{\frac{k+l}{2}\frac{m}{2}}}{1+|f(j)-i|}\right]^{s-1} \end{aligned} \tag{63}$$

and according to lemma 7.1

$$\begin{aligned} \sum_{ij} |a_j|^s |b_i|^s \left[\frac{2^{\frac{k+l}{2}\frac{m}{2}}}{1+|f(j)-i|}\right]^{s-1} \lesssim& \|a^s\|_r \|b^s\|_r \|G_{\frac{k+l}{2}}^{s-1}\|_{\frac{r'}{2}} \\ \overset{(62)}{\lesssim}& \|a^s\|_r \|b^s\|_r 2^{\frac{k+l}{2}m\left(1-\frac{s}{q'}\right)} \\ \overset{(59)}{=}& \|a\|_{q'}^s \|b\|_{q'}^s 2^{\frac{k+l}{2}m\left(1-\frac{s}{q'}\right)}. \end{aligned}$$

Putting this into (63) yields the claim, when we insert the expression $2^{-\frac{|k-l|}{2}(m-1)(s-1)}$, which is of constant order in case of $|k-l| \lesssim 1$. ☐

Eventually we can complete the proof of theorem 2.2. Let me remind you of its statement:

THEOREM
Let $p' > m+1$, $\frac{3}{p'} < \frac{1}{q} < \frac{1}{2} + \frac{1}{p'}$, *then*

$$\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'} \lesssim \delta^{\frac{1}{q}} \| \sum_{\alpha=(k,j,n)} a_\alpha ((2^k\text{-}1)\gamma)^{\frac{1}{q} - \frac{m+1}{p'}} \phi_\alpha \|_{q'} \tag{64}$$

holds for all $\delta > 0$ *and for all sequences* a_α.

Proof: Since $2 < p' < \infty$, we find $s \in (1, \infty)$ such that $\frac{1}{s} + \frac{2}{p'} = 1$. (65)
Since even $4 < p'$, $\frac{p'}{2} > 2$ holds, thus we can apply Youngs inequality:

$$\begin{aligned}
\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'}^{2s} &= \| \sum_{\alpha,\mu} a_\alpha a_\mu \hat{\phi}_\alpha \hat{\phi}_\mu \|_{\frac{p'}{2}}^{s} \\
&= \| \mathfrak{F} \Big(\sum_{\alpha,\mu} a_\alpha a_\mu \phi_\alpha * \phi_\mu \Big) \|_{\frac{p'}{2}}^{s} \\
&\overset{Young}{\leq} \| \sum_{\alpha,\mu} a_\alpha a_\mu \phi_\alpha * \phi_\mu \|_{s}^{s} \\
&= \int \Big| \sum_{\alpha,\mu} a_\alpha a_\mu \phi_\alpha * \phi_\mu \Big|^{s} \,\mathrm{d}x.
\end{aligned}$$

Now we exploit the estimate from chapter 4 of the overlap of the supports of the functions $\phi_\alpha * \phi_\mu$:

$$\begin{aligned}
\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'}^{2s} &\leq \int \Big| \sum_{\alpha,\mu} a_\alpha a_\mu \phi_\alpha * \phi_\mu \Big|^{s} \,\mathrm{d}x. \\
&\overset{\text{corollary } 4.6}{\lesssim} \beta^{1-s} \sum_{\alpha,\mu} |a_\alpha|^s |a_\mu|^s \int |\phi_\alpha * \phi_\mu|^s \,\mathrm{d}x \\
&\overset{\text{lemma } 6.2}{\lesssim} \beta^2 (\delta\gamma)^{s+1} \sum_{klnp} \|a_{kn}\|_{q'}^{s} \|a_{lp}\|_{q'}^{s} 2^{\frac{k+l}{2}(1+s-sm+m\left(1-\frac{s}{q'}\right))} 2^{-\frac{|k-l|}{2}(m-1)(s-1)},
\end{aligned}$$

where $a_{kn} = (a_{kjn})_{j \in I_k}$, $a_{lp} = (a_{lip})_{i \in I_l}$. Notice that for fixed k Hölder implies

$$\begin{aligned}
\sum_{n} \|a_{kn}\|_{q'}^{s} &\leq \left(\sum_{n} \|a_{kn}\|_{q'}^{q'} \right)^{\frac{s}{q'}} \left(\sum_{n=\frac{1}{\beta}}^{\frac{2}{\beta}-1} 1 \right)^{1-\frac{s}{q'}} \\
&= \|a_k\|_{q'}^{s} \left(\frac{1}{\beta} \right)^{1-\frac{s}{q'}} = \|a_k\|_{q'}^{s} \beta^{\frac{s}{q'}-1},
\end{aligned} \tag{66}$$

where $a_k = (a_{kjn})_{j,n}$. Implementing this in our equations yields

$$\begin{aligned}
\|\sum_\alpha a_\alpha \hat{\phi}_\alpha\|_{p'}^{2s} \lesssim & \beta^{2(\frac{s}{q'}-1)}\beta^2(\delta\gamma)^{(s+1)}\sum_{k,l}\|a_k\|_{q'}^s\|a_l\|_{q'}^s 2^{\frac{k+l}{2}(1+s-sm+m\left(1-\frac{s}{q'}\right))}2^{-|k-l|\frac{(m-1)(s-1)}{2}} \\
= & \beta^{\frac{2s}{q'}}(\delta\gamma)^{(s+1)}\sum_{k,l}\|a_k\|_{q'}^s\|a_l\|_{q'}^s 2^{\frac{k+l}{2}[s(1-m)+m+1-m\frac{s}{q'}]}2^{-|k-l|\frac{(m-1)(s-1)}{2}} \\
= & \beta^{\frac{2s}{q'}}(\delta\gamma)^{(s+1)}\sum_{k,l}\|a_k 2^{k[\frac{1-m}{2}+\frac{m+1}{2s}-\frac{m}{2q'}]}\|_{q'}^s\|a_l 2^{l[\frac{1-m}{2}+\frac{m+1}{2s}-\frac{m}{2q'}]}\|_{q'}^s 2^{-|k-l|\frac{(m-1)(s-1)}{2}}.
\end{aligned} \tag{67}$$

We again apply lemma 7.1 for $r = \frac{q'}{s}$ (recall $r < 2$, cf.(60)):

$$\begin{aligned}
\|\sum_\alpha a_\alpha \hat{\phi}_\alpha\|_{p'}^{2s} \lesssim \quad & \beta^{\frac{2s}{q'}}(\delta\gamma)^{s+1}\left(\sum_k \|a_k 2^{k[\frac{1-m}{2}+\frac{m+1}{2s}-\frac{m}{2q'}]}\|_{q'}^{q'}\right)^{\frac{2s}{q'}} \\
= \quad & (\delta\gamma)^{s+1-\frac{2s}{q'}}\gamma^{-\left(\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}\right)2s}\left(\sum_k \|a_k(2^k\gamma)^{\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}}\|_{q'}^{q'}2^{k(1-\frac{m}{2})}\gamma\beta\delta\right)^{\frac{2s}{q'}} \\
\overset{\delta=\gamma^m}{=} \quad & \gamma^{(m+1)\left(s+1-\frac{2s}{q'}\right)-\left(\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}\right)2s}\left(\sum_k \|a_k(2^k\gamma)^{\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}}\|_{q'}^{q'}2^{k(1-\frac{m}{2})}\gamma\beta\delta\right)^{\frac{2s}{q'}}.
\end{aligned} \tag{68}$$

We determine the exponents: It holds

$$\begin{aligned}
&(m+1)\left(s+1-\frac{2s}{q'}\right)-\left(\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}\right)2s \\
=&2s\left[(m+1)\left(\frac{1}{2}+\frac{1}{2s}-\frac{1}{q'}\right)-\frac{1-m}{2}-\frac{m+1}{2s}+\frac{1}{q'}\right] \\
=&2s\left[\frac{m+1}{2}-\frac{m}{q'}-\frac{1-m}{2}\right] \\
=&2s\,m\left[1-\frac{1}{q'}\right]=\frac{2sm}{q},
\end{aligned} \tag{69}$$

and

$$\frac{1-m}{2}+\frac{m+1}{2s}-\frac{1}{q'}=\frac{1}{q}-\frac{m+1}{2}+\frac{m+1}{2s}=\frac{1}{q}-\frac{m+1}{2}\left(1-\frac{1}{s}\right)\overset{(65)}{=}\frac{1}{q}-\frac{m+1}{p'}. \tag{70}$$

A further computation shows

$$\int \phi_\alpha^{q'}\,\mathrm{d}x \approx |\Gamma_\alpha^\delta| = |\Gamma_{kjn}^\delta| \approx \gamma_k\beta\delta = 2^{k(1-\frac{m}{2})}\gamma\beta\delta, \tag{71}$$

thus (68) translates into

$$\begin{aligned}
\| \sum_{\alpha} a_\alpha \hat{\phi}_\alpha \|_{p'}^{2s} \lesssim & \gamma^{m\frac{2s}{q}} \left(\sum_k \| a_k (2^k \gamma)^{\frac{1}{q} - \frac{m+1}{p'}} \|_{q'}^{q'} \int \phi_\alpha^{q'} \ \mathrm{d}x \right)^{\frac{2s}{q'}} \\
= & \delta^{\frac{2s}{q}} \left(\int \sum_{\alpha} \left| a_\alpha (2^k \gamma)^{\frac{1}{q} - \frac{m+1}{p'}} \phi_\alpha \right|^{q'} \ \mathrm{d}x \right)^{\frac{2s}{q'}} \\
\approx & \left(\delta^{\frac{1}{q}} \| \sum_{\alpha} a_\alpha ((2^k\text{-}1)\gamma)^{\frac{1}{q} - \frac{m+1}{p'}} \phi_\alpha \|_{q'} \right)^{2s},
\end{aligned}$$

completing the proof. □

7 Appendix: Auxiliary lemmas

7.1 Classical lemmas

Lemma 7.1
Let $1 \leq r < 2$, *and* $f : \mathbb{N} \to \mathbb{N}$ *„almost“ injectiv, i.e. there exists a constant* $C > 0$ *such that* $\forall l : |\{k \in \mathbb{N} : f(k) = l\}| \leq C$. (72)
Then

$$\left|\sum_{k,l} a_k G_{f(k)-l} b_l\right| \leq C^{\frac{1}{r'}} \|a\|_r \|G\|_{\frac{r'}{2},\infty} \|b\|_r$$

holds for all sequences $a, b \in \ell_r$, $G \in \ell_{\frac{r'}{2},\infty}$.

Proof: At first, Hölder's inequality gives

$$\left|\sum_{k,l} a_k G_{f(k)-l} b_l\right| \leq \|a\|_r \|(G * b)(f(\cdot))\|_{r'}.$$

We further observe

$$\sum_k |G * b|^{r'}(f(k)) \underset{(72)}{\leq} C \sum_l |G * b|^{r'}(l).$$

Since $r < 2$, we have $r' > 2$, i.e. $\frac{r'}{2} > 1$. Hence we apply Young's inequality with parameters $1 + \frac{1}{r'} = \frac{2}{r'} + \frac{1}{r}$ to obtain

$$\|G * b\|_{r'} \leq \|G\|_{\frac{r'}{2},\infty} \|b\|_r.$$

Altogether, this provides the desired estimate. ☐

Lemma 7.2 (Hölder's inequality in Lorentz spaces)
Let $\frac{1}{p_i} = \frac{1}{q_i} + \frac{1}{r_i}$, $i = 1, 2$. *Then*

$$\|fg\|_{p_1,p_2} \leq \|f\|_{q_1,q_2} \|g\|_{r_1,r_2}.$$

Proof: [G] ☐

Lemma 7.3 (Brunn-Minkowski)
Let K_0, K_1 *be compact and convex subsets of* $\mathbb{R}^n$ *and* $0 \leq t \leq 1$. *Then*

$$(1-t)\,|K_0|^{\frac{1}{n}} + t\,|K_1|^{\frac{1}{n}} \leq |(1-t)K_0 + tK_1|^{\frac{1}{n}}.$$

Proof: Compare [GW], theorem 6.2, p. 57. ☐

7.2 More special lemmas

Lemma 7.4
Let K and L be two convex bodies in $\mathbb{R}^n$ symmmetric with respect to the origin (i.e. $K=-K$). Then $\sup_{z\in\mathbb{R}^n} |K\cap(z+L)| = |K\cap L|$ holds for all $z\in\mathbb{R}^n$.

It is a simple observation that both conditions on K and L are crucial.

Proof: Let z be some point in $\mathbb{R}^n$. We have to show that $|K\cap(z+L)|\leq|K\cap L|$. Therefore we introduce the sets $A=(L+z)\cap K$ and $B=(L-z)\cap K$. The proof will be divided in three steps:

(i) $\frac{A+B}{2}\subset K\cap L$

(ii) $|A|=|B|$

(iii) $|A|\leq\left|\frac{A+B}{2}\right|$. □

Notice that (iii) and (i) immediatly imply what we claimed.
Proof of (i): On the one hand, since $A\subset K$ and $B\subset K$, we have $\frac{A+B}{2}\subset\frac{K+K}{2}\overset{K \text{ convex}}{=}K$. On the other hand, since $A\subset L+z$, $B\subset L-z$, we have $\frac{A+B}{2}\subset\frac{L+z+L-z}{2}\overset{L \text{ convex}}{=}L$.
Proof of (ii): This is clear since $A=-B$ by the symmetry condition on K and L.
Proof of (iii): We apply the inequality of Brunn-Minkowski:

$$\begin{aligned} |A| &= \left(\frac{1}{2}|A|^{\frac{1}{n}}+\frac{1}{2}|A|^{\frac{1}{n}}\right)^n \\ &\overset{(ii)}{=} \left(\frac{1}{2}|A|^{\frac{1}{n}}+\frac{1}{2}|B|^{\frac{1}{n}}\right)^n \\ &\underset{\text{Minkowski}}{\overset{\text{Brunn-}}{\leq}} \left|\frac{A+B}{2}\right|. \end{aligned}$$

Lemma 7.5
Let X be a normed vectorspace and $U_1,U_2,V_1,V_2\subset X$ with the property $U_1+V_1\cap U_2+V_2\neq\emptyset$. Furthermore let P be a linear functional on X and let $x_i\in U_i$, $a=P(x_2)-P(x_1)$ and $y_i\in V_i$, $b=P(y_2)-P(y_1)$. Then we have

$$a+b\leq\sum_{i=1,2} diam(P(U_i))+diam(P(V_i)). \tag{73}$$

Proof: Choose some $\xi\in U_1+V_1\cap U_2+V_2$. There exists $u_i\in U_i$, $v_i\in V_i$, $i=1,2$ with $u_1+v_1=\xi=u_2+v_2$. It follows

$$\begin{aligned} a+b =& P(x_2-x_1)+P(y_2-y_1) \\ =& P(x_2-x_1)+P(y_2-y_1)+P(u_1+v_1)-P(u_2+v_2) \\ =& P(x_2-u_2)+P(y_2-v_2)+P(u_1-x_1)+P(v_1-y_1) \\ \leq& diam(P(U_2))+diam(P(V_2))+diam(P(U_1))+diam(P(V_1)), \end{aligned}$$

completing the proof. □

Lemma 7.6

Let $I = \{0,\dots,n\}, J = \{0,\dots,m\} \subset \mathbb{N}$ *and let* $a = (a_i)_{i\in I} \in \mathbb{R}^I$, $b = (b_j)_{j\in J} \in \mathbb{R}^J$ *be two increasing, finite sequences such that*

$$\begin{array}{ll} C^{-1} \leq \ a_{i+1} - a_i \leq C & \textit{für alle } i \in I \\ C^{-1} \leq \ b_{j+1} - b_j \leq C & \textit{für alle } j \in J. \end{array} \tag{74}$$

Moreover we assume that $b_m - b_0 \leq a_n - a_0$. (75)

Then there exists a function $f : J \to I$ *such that*

$$\begin{array}{ll} (i) & |a_i - b_j| \geq \frac{1}{2}|a_i - a_{f(j)}| \\ (ii) & \forall i \in I : |\{j : f(j) = i\}| \leq 4C^2 + 2. \end{array}$$

The trivial case $I = J$, $a = b$ can of course be solved by $f = id$. In the general setting, we in some sense had to replace every b_j by some $a_{f(j)}$, leaving the distances to other points (almost) unchanged. Condition (ii) can be read as a weakening of injectivity.

Proof:

Without loss of generality, we may assume $b_m \leq a_n$. (76)

To be more precise, if $b_m > a_n$ would hold, we would consider $\bar{a}_i = -a_{n-i}$ and $\bar{b}_j = -b_{m-j}$, which also fulfill the requirements of the lemma. Then

$$\bar{b}_m = -b_0 \underset{(75)}{\leq} a_n - a_0 - b_m < -a_0 = \bar{a}_n.$$

So, if we would find a function $\bar{f}$ appropriate to $\bar{a}$, $\bar{b}$ in the sense of (i) and (ii), $f(j) := n - \bar{f}(m-j)$ would be a solution appropriate to a and b since

$$|a_i - b_j| = |\bar{a}_{n-i} - \bar{b}_{m-j}| \geq \frac{1}{2}|\bar{a}_{n-i} - \bar{a}_{\bar{f}(m-j)}| = \frac{1}{2}|a_i - a_{n-\bar{f}(m-j)}|.$$

Furthermore, we may also assume $a_0 \leq b_m$ without loss of generality. (77)

In the case $a_0 > b_m$, we would introduce $\bar{a}_i = a_i - a_0 + b_m < a_i$, such that $\bar{a}_0 = b_m \leq b_m$. If f is a function associated to $\bar{a}$ and b as required, then

$$|b_j - a_i| \underset{b_m<a_0}{=} a_i - b_j > \bar{a}_i - b_j \underset{b_m=\bar{a}_0}{=} |b_j - \bar{a}_i| \geq \frac{1}{2}|\bar{a}_{f(j)} - \bar{a}_i| = \frac{1}{2}|a_{f(j)} - a_i|.$$

It would be helpfull for the construction of f to assign an i to every given j in a way that b_j is close to a_i. Nevertheless, if the sequences are somehow shifted against each others (for instance, $b_0 \ll a_0$), there might be no a_i „close“ to b_j. If we would always choose the closest a_i, this would hurt condition (ii).

Therefore we mirror $a_i's$ at a_0, to ensure that for every b_j there is a (maybe mirrored) point a_i nearby.



Fig. 9: Setting of the sequences a and b

Let $\bar{I} = I \cup (-I)$. Define the continuation of a on $\bar{I}$ by $a_{-i} = 2a_0 - a_i$, $i \in I$. We then find a function $g : J \to \bar{I}$ such that $|a_{g(j)} - b_j| = \min_{i \in \bar{I}} |a_i - b_j|$. The claim is that $f = |g|$ is asolution of our problem.

Checking condition (i):

Case 1: $g(j) \geq 0$. Here

$$|a_i - a_{f(j)}| = |a_i - a_{g(j)}| \leq |a_i - b_j| + |b_j - a_{g(j)}| \underset{\text{minimality of } g}{\leq} 2|a_i - b_j|.$$

Case 2: $g(j) < 0$. Here

$$\begin{aligned}|a_i - a_{f(j)}| = |a_i - a_{-g(j)}| &= |a_i - 2a_0 + a_{g(j)}| \\ &\leq |a_i - a_0| + |a_0 - a_{g(j)}| \underset{\text{monotony}}{=} a_i - a_0 + a_0 - a_{g(j)} = |a_i - a_{g(j)}|,\end{aligned}$$

and we proceed as in case 1.

Checking condition (ii):

Obviously, it is sufficient to show that $|\{j : g(j) = i\}| \leq 2C^2 + 1$ holds for every $i \in \bar{I}$. Thus let $g(j) = i$.

We claim that $|a_i - b_j| \leq C$ and check this:

Case 1: $a_i < b_j$. Would $i = n$, then $b_j \leq b_m \underset{(76)}{\leq} a_n = a_i < b_j$, hence $i < n$. Thus a_{i+1} is well-defined and $b_j < a_{i+1}$ holds according to the minimality in the choise of g. We conclude $|a_i - b_j| = b_j - a_i \leq a_{i+1} - a_i \leq C$.

Case 2: $a_i > b_j$. Would $i = -n$, then

$$b_j < a_i = a_{-n} = 2a_0 - a_n \underset{(77)}{\leq} b_m - (a_n - a_0) \underset{(75)}{\leq} b_0 \leq b_j.$$

Hence we have $i > -n$, thus a_{i-1} is well-defined and $b_j > a_{i-1}$ holds. We finish as in case 1.

Case 3: $a_i = b_j$. This case is trivial.

If additionally $g(j') = i$, then $|a_i - b_j| \leq C$ and $|a_i - b_{j'}| \leq C$. Now we utilise (74). Therefrom we get by induction

$$\frac{1}{C}|j - j'| \leq |b_j - b_{j'}| \leq |b_j - a_i| + |a_i - b_{j'}| \leq 2C,$$

i.e.

$$|j - j'| \leq 2C^2,$$

and thus condition (ii). □

REMARK

If assumption (75) *is not fulfilled, a comparable result can be obtained by interchanging the roles of I and J or a and b respectively. Actually, the lemma remains also essentially valid if these quantities are* not *interchanged. Therefore we just need some further mirrorings, if necessary also at the „upper“ end a_n. This would enlarge the bound $4C^2 + 2$ from (ii) depending on the relation between n and m. Nevertheless, the notation would quickly become confusing, thus I omit such a version of the lemma.*

References

[ACK] Arkhipov, Karatsuba, Chubarikov, Trigonometric integrals, Mathematics of the USSR Izvestiya 15 No2 (1980), 211-239

[Ba1] Barcelo Taberner, On the restriction of the fourier transform to a conical surface, Transactions of the american mathematical Society, 292 No1 (1985)

[Ba2] Barcelo Taberner, The restriction of the Fourier transform to some curves and surfaces, Studia Mathematica 84 No1 (1986), 39–69

[Bu] Burwell, Asymptotic expansions of generalized hypergeometric functions, Proceedings of the London Mathematical Society (2) 22, 57-72

[DM] Drury, Marshall, Fourier restriction theorems for curves with affine and Euclidean arclengths, Mathematical Proceedings of the Cambridge Philosophical Society 97 (1985), 111-125

[G] Grafakos, Classical and Modern Fourier Analysis, Pearson Education Inc., 2004

[GW] Gruber, Wills, Handbook of convex geometry, North-Holland (1993)

[St] Stein, Harmonic Analysis, Princeton University Press (1993)

[StSh] Stein, Shakarchi, Fourier Analysis, Princeton Lectures in Analysis (2003)

[T] Tao, Recent progress on the Restriction conjecture, arXiv:math/0311181v1 [math.CA] 12 Nov 2003